



PLAINTIFF'S
 EXHIBIT
 3

Drug Facts

Active ingredients (in each tablet) Purpose
Loratadine 10 mg..... Antihistamine
Pseudoephedrine sulfate 240 mg..... Nasal decongestant

Uses

- temporarily relieves those symptoms due to hay fever or other upper respiratory allergies:
 - sneezing
 - itching of the nose or throat
 - runny nose
 - itchy, watery eyes
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores free breathing through the nose

Warnings

- Do not use
- If you have ever had an allergic reaction to this product or any of its ingredients.
 - If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- Ask a doctor before use if you have
- heart disease
 - high blood pressure
 - trouble urinating due to an enlarged prostate gland
 - liver or kidney disease. Your doctor should determine if you need a different dose.
 - thyroid disease
 - diabetes

Drug Facts (continued)

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if:

- an allergic reaction to this product occurs.
 - Seek medical help right away.
 - symptoms do not improve within 7 days or are accompanied by a fever.
 - irritation, dizziness or sleeplessness occurs.
- If pregnant or breast-feeding, ask a health professional before use.
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not divide, crush, chew or dissolve the tablet
- adults and children 12 years and over: 1 tablet daily with a full glass of water; not more than 1 tablet in 24 hours
- children under 12 years of age: ask a doctor
- consumers with liver or kidney disease: ask a doctor

Other information

- do not use if blister unit is broken or torn
- store between 20° to 25°C (68° to 77°F)
- protect from light and store in a dry place

Inactive ingredients black iron oxide, candellila wax powder, colloidal silicon dioxide, glyceryl monostearate, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, polyisobutyl 20, polyethylene glycol, sodium lauryl sulfate, talc, titanium dioxide

Drug Facts (continued)
(Questions or comments? 1-800-774-2578)

This product is not manufactured or distributed by Schering-Plough HealthCare Products, Inc., distributor of Claritin, P.O. Box 1400, Kenilworth, NJ 07033. © 1998 Schering-Plough HealthCare Products, Inc. All rights reserved. For Dosage and Administration information, see the accompanying Patient Information Statement. For more information, call 1-800-774-2578. WML/evs/leo.com

