

**EXHIBIT B**  
**Part 2 of 2**

4,703,008

21

with a 1 M NaCl —1% SDS buffer for 55° C. for 4 hours or more, after which the buffer was removed. Hybridization and post-hybridization washings were carried out as described in Example 3, Part B. Both the mixture of 128 20-mer probes designated EPV and the mixture of 128 17-mer probes of Table III (designated the EPQ mixture) were employed. Hybridization was carried out at 48° C. using the EPV probe mixture. EPQ probe mixture hybridization was carried out at 46° C. —4 degrees below the lowest calculated Td for members of the mixture. Removal of the hybridized probe for rehybridization was accomplished by boiling with 1×SSC —0.1% SDS for two minutes. Autoradiography of the filters revealed three positive clones (reactive with both probe mixtures) among the 1,500,000 phage plaques screened. Verification of the positive clones as being EPO-encoding was obtained through DNA sequencing and electron micrographic visualization of heteroduplex formation with the monkey cDNA of Example 3. This procedure also gave evidence of multiple introns in the genomic DNA sequence.

#### EXAMPLE 5

Nucleotide sequence analysis of one of the positive clones (designated  $\lambda$ AHE1) was carried out and results obtained to date are set out in FIG. 6.

In FIG. 6, The initial continuous DNA sequence designates a top strand of 620 bases in what is apparently an untranslated sequence immediately preceding a translated portion of the human EPO gene. More specifically, the sequence appears to comprise the 5' end of the gene which leads up to a translated DNA region coding for the first four amino acids (—27 through —24) of a leader sequence ("presequence"). Four base pairs in the sequence prior to that encoding the beginning of the leader have not yet been unambiguously determined and are therefore designated by an "X". There then follows an intron of about 639 base pairs (439 base pairs of which have been sequenced and the remaining 200 base pairs of which are designated "I.S.") and immediately preceding a codon for glutamine which has been designated as residue —23 of the translated polypeptide. The exon sequence immediately following is seen to code for amino acid residues through an alanine residue (designated as the +1 residue of the amino acid sequence of mature human EPO) to the codon specifying threonine at position +26, whereupon there follows a second intron consisting of 256 bases as specifically designated. Following this intron is an exon sequence for amino acid residues 27 through 55 and thereafter a third intron comprising 612 base pairs commences. The subsequent exon codes for residues 56 through 115 of human EPO and there then commences a fourth intron of 134 bases as specified. Following the fourth intron is an exon coding for residue Nos. 116 through 166 and a "stop" codon (TGA). Finally, Table VI identifies a sequence of 568 base pairs in what appears to be an untranslated 3' region of the human EPO gene, two base pairs of which ("X") have not yet been unambiguously sequenced.

FIG. 6 thus serves to identify the primary structural conformation (amino acid sequence) of mature human EPO as including 166 specified amino acid residues (estimated M.W. = 18,399). Also revealed in the Figure is the DNA sequence coding for a 27 residue leader sequence along with 5' and 3' DNA sequences which may be significant to promoter/operator functions of the human gene operon. Sites for potential glycosyla-

22

tion of the mature human EPO polypeptide are designated in the Figure by asterisks. It is worthy of note that the specific amino acid sequence of FIG. 6 likely constitutes that of a naturally occurring allelic form of human erythropoietin. Support for this position is found in the results of continued efforts at sequencing of urinary isolates of human erythropoietin which provided the finding that a significant number of erythropoietin molecules therein have a methionine at residue 126 as opposed to a serine as shown in the Figure.

FIG. 9 illustrates the extent of polypeptide sequence homology between human and monkey EPO. In the upper continuous line of the Figure, single letter designations are employed to represent the deduced translated polypeptide sequences of human EPO commencing with residue —27 and the lower continuous line shows the deduced polypeptide sequence of monkey EPO commencing at assigned residue number —27. Asterisks are employed to highlight the sequence homologies. It should be noted that the deduced human and monkey EPO sequences reveal an "additional" lysine (K) residue at (human) position 116. Cross-reference to FIG. 6 indicates that this residue is at the margin of a putative mRNA splice junction in the genomic sequence. Presence of the lysine residue in the human polypeptide sequence was further verified by sequencing of a cDNA human sequence clone prepared from mRNA isolated from COS-1 cells transformed with the human genomic DNA in Example 7, infra.

#### EXAMPLE 6

The expression system selected for initial attempts at microbial synthesis of isolatable quantities of EPO polypeptide material coded for by the monkey cDNA provided by the procedures of Example 3 was one involving mammalian host cells (i.e., COS-1 cells, A.T.C.C. No. CRL-1650). The cells were transfected with a "shuttle" vector capable of autonomous replication in *E. coli* host (by virtue of the presence of pBR322-derived DNA) and the mammalian hosts (by virtue of the presence of SV40 virus-derived DNA).

More specifically, an expression vector was constructed according to the following procedures. The plasmid clone 83 provided in Example 3 was amplified in *E. coli* and the approximately 1.4 kb monkey EPO-encoding DNA was isolated by EcoRI and HindIII digestion. Separately isolated was an approximately 4.0 kb, HindIII/SalI fragment from pBR322. An approximately 30 bp, EcoRI/SalI "linker" fragment was obtained from M13mp10 RF DNA (P and L Laboratories). This linker included, in series, an EcoRI sticky end, followed by SstI, SmaI, BamHI and XbaI recognition sites and a SalI sticky end. The above three fragments were ligated to provide an approximately 5.4 kb intermediate plasmid ("pERS") wherein the EPO DNA was flanked on one side by a "bank" of useful restriction endonuclease recognition sites. pERS was then digested with HindIII and SalI to yield the EPO DNA and the EcoRI to SalI (M13mp10) linker. The 1.4 kb fragment was ligated with an approximately 4.0 kb BamHI/SalI of pBR322 and another M13mp10 HindIII/BamHI RF fragment linker also having approximately 30 bp. The M13 linker fragment was characterized by a HindIII sticky end, followed by PstI, SalI, XbaI recognition sites and a BamHI sticky end. The ligation product was, again, a useful intermediate plasmid ("pBR-EPO") including the EPO DNA flanked on both sides by banks of restriction site.

4,703,008

23

The vector chosen for expression of the EPO DNA in COS-1 cells ("pDSVL1") had previously been constructed to allow for selection and autonomous replication in *E. coli*. These characteristics are provided by the origin of replication and Ampicillin resistance gene DNA sequences present in the region spanning nucleotides 2448 through 4362 of pBR322. This sequence was structurally modified by the addition of a linker providing a HindIII recognition immediately adjacent nucleotide 2448 prior to incorporation into the vector. Among the selected vector's other useful properties was the capacity to autonomously replicate in COS-1 cells and the presence of a viral promoter sequence functional in mammalian cells. These characteristics are provided by the origin of replication DNA sequence and "late gene" viral promoter DNA sequence present in the 342 bp sequence spanning nucleotide numbers 5171 through 270 of the SV40 genome. A unique restriction site (BamHI) was provided in the vector and immediately adjacent the viral promoter sequence through use of a commercially available linker sequence (Collaborative Research). Also incorporated in the vector was a 237 base pair sequence (derived as nucleotide numbers 2553 through 2770 of SV40) containing the "late gene" viral mRNA polyadenylation signal (commonly referred to as a transcription terminator). This fragment was positioned in the vector in the proper orientation vis-a-vis the "late gene" viral promoter via the unique BamHI site. Also present in the vector was another mammalian gene at a location not material to potential transcription of a gene inserted at the unique BamHI site, between the viral promoter and terminator sequences. [The mammalian gene comprised an approximately 2,500 bp mouse dihydrofolate reductase (DMFR) mini-gene isolated from plasmid pMG-1 as in Gasser, et al., *P.N.A.S. (U.S.A.)*, 79, pp. 6522-6526, (1982).] Again, the major operative components of plasmid pDSVL1 comprise nucleotides 2448 through 4362 of pBR322 along with nucleotides 5171 through 270 (342 bp) and 2553 through 2770 (237 bp) of SV40 DNA.

Following procedures described, e.g., in Maniatis, et al., *supra*, the EPO-encoding DNA was isolated from plasmid pBR-EPO as a BamHI fragment and ligated into plasmid pDSVL1 cut with BamHI. Restriction enzyme analysis was employed to confirm insertion of the EPO gene in the correct orientation in two of the resulting cloned vectors (duplicate vectors H and L). See FIG. 2, illustrating plasmid pDSVL-MkE. Vectors with EPO genes in the wrong orientation (vectors F, X and G) were saved for use as negative controls in transfection experiments designed to determine EPO expression levels in hosts transformed with vectors having EPO DNA in the correct orientation.

Vectors H, L, F, X and G were combined with carrier DNA (mouse liver and spleen DNA) were employed to transfect duplicate 60 mm plates by calcium phosphate microprecipitate methods. Duplicate 60 mm plates were also transfected with carrier DNA as a "mock" transformation negative control. After five days all culture media were tested for the presence of polypeptides possessing the immunological properties of naturally-occurring EPO.

#### EXAMPLE 7

##### A. Initial EPO Expression System Involving COS-1 Cells

The system selected for initial attempts at microbial synthesis of isolatable quantities of human EPO poly-

24

peptide material coded for by the human genomic DNA EPO clone, also involved expression in mammalian host cells (i.e., COS-1 cells, A.T.C.C. No. CRL-1650). The human EPO gene was first sub-cloned into a "shuttle" vector which is capable of autonomous replication in both *E. coli* hosts (by virtue of the presence of pBR322 derived DNA) and in the mammalian cell line COS-1 (by virtue of the presence of SV40 virus derived DNA). The shuttle vector, containing the EPO gene, was then transfected into COS-1 cells. EPO polypeptide material was produced in the transfected cells and secreted into the cell culture media.

More specifically, an expression vector was constructed according to the following procedures. DNA isolated from lambda clone  $\lambda$ hE1, containing the human genomic EPO gene, was digested with BamHI and HindIII restriction endonucleases, and a 5.6 kb DNA fragment known to contain the entire EPO gene was isolated. This fragment was mixed and ligated with the bacterial plasmid pUC8 (Bethesda Research Laboratories, Inc.) which had been similarly digested, creating the intermediate plasmid "pUC8-HuE", providing a convenient source of this restriction fragment.

The vector chosen for expression of the EPO DNA in COS-1 cells (pSV4SEt) had previously been constructed. Plasmid pSV4SEt contained DNA sequences allowing selection and autonomous replication in *E. coli*. These characteristics are provided by the origin of replication and Ampicillin resistance gene DNA sequences present in the region spanning nucleotides 2448 through 4362 of the bacterial plasmid pBR322. This sequence was structurally modified by the addition of a linker providing a HindIII recognition site immediately adjacent to nucleotide 2448. Plasmid pSV4SEt was also capable of autonomous replication in COS-1 cells. This characteristic was provided by a 342 bp fragment containing the SV40 virus origin of replication (nucleotide numbers 5171 through 270). This fragment had been modified by the addition of a linker providing an EcoRI recognition site adjacent to nucleotide 270 and a linker providing a Sall recognition site adjacent nucleotide 5171. A 1061 bp fragment of SV40 was also present in this vector (nucleotide numbers 1711 through 2772 plus a linker providing a Sall recognition site next to nucleotide number 2772). Within this fragment was an unique BamHI recognition sequence. In summary, plasmid pSV4SEt contained unique BamHI and HindIII recognition sites, allowing insertion of the human EPO gene, sequences allowing replication and selection in *E. coli*, and sequences allowing replication in COS-1 cells.

In order to insert the EPO gene into pSV4SEt, plasmid pUC8-HuE was digested with BamHI and HindIII restriction endonucleases and the 5.6 kb EPO encoding DNA fragment isolated. pSV4SEt was also digested with BamHI and HindIII and the major 2513 bp fragment isolated (preserving all necessary functions). These fragments were mixed and ligated, creating the final vector "pSVgHuEPO". (See, FIG. 3.) This vector was propagated in *E. coli* and vector DNA isolated. Restriction enzyme analysis was employed to confirm insertion of the EPO gene.

Plasmid pSVgHuEPO DNA was used to express human EPO polypeptide material in COS-1 cells. More specifically, pSVgHuEPO DNA was combined with carrier DNA and transfected into triplicate 60 mm plates of COS-1 cells. As a control, carrier DNA alone was also transfected into COS-1 cells. Cell culture media were sampled five and seven days later and tested

4,703,008

25

for the presence of polypeptides possessing the immunological properties of naturally occurring human EPO.

#### B. Second EPO Expression System Involving COS-1 Cells

Still another system was designed to provide improved production of human EPO polypeptide material coded by the human genomic DNA EPO clone in COS-1 cells (A.T.C.C. No. CRL-1650).

In the immediately preceding system, EPO was expressed in COS-1 cells using its own promoter which is within the 5.6 Kb BamHI to HindIII restriction fragment. In the following construction, the EPO gene is altered so that it is expressed using the SV40 late promoter.

More specifically, the cloned 5.6 Kb BamHI to HindIII genomic human EPO restriction fragment was modified by the following procedures. Plasmid pUC8-HuE, as described above, was cleaved with BamHI and with BstEII restriction endonucleases. BstEII cleaves within the 5.6 Kb EPO gene at a position which is 44 base pairs 5' to the initiating ATG coding for the prepeptide and approximately 680 base pairs 3' to the HindIII restriction site. The approximately 4900 base pair fragment was isolated. A synthetic linker DNA fragment, containing SalI and BstEII sticky ends and an internal BamHI recognition site was synthesized and purified. The two fragments were mixed and ligated with plasmid pBR322 which had been cut with SalI and BamHI to produce the intermediate plasmid pBRgHE. The genomic human EPO gene can be isolated therefrom as a 4900 base pair BamHI digestion fragment carrying the complete structural gene with a single ATG 44 base pairs 3' to BamHI site adjacent the amino terminal coding region.

This fragment was isolated and inserted as a BamHI fragment into BamHI cleaved expression vector plasmid pDSVL1 (described in Example 6). The resulting plasmid, pSVLgHuEPO, as illustrated in FIG. 4, was used to express EPO polypeptide material from COS-1 cells, as described in Examples 6 and 7A.

#### EXAMPLE 8

Culture media from growth of the six transfected COS-1 cultures of Example 6 were analyzed by radio-immunoassay according to the procedures set forth in Example 2, Part B. Each sample was assayed at 250, 125, 50, and 25 microliter aliquot levels. Supernatants from growth of cells mock transfected or transfected with vectors having incorrect EPO gene orientation were unambiguously negative for EPO immunoreactivity. For each sample of the two supernatants derived from growth of COS-1 cells transfected with vectors (H and L) having the EPO DNA in the correct orientation, the % inhibition of <sup>125</sup>I-EPO binding to antibody ranged from 72 to 88%, which places all values at the top of the standard curve. The exact concentration of EPO in the culture supernatant could not then reliably be estimated. A quite conservative estimate of 300 mU/ml was made, however, from the value calculation of the largest aliquot size (250 microliter).

A representative culture fluid according to Example 6 and five and seven day culture fluids obtained according to Example 7A were tested in the RIA in order to compare activity of recombinant monkey and human EPO materials to a naturally-occurring human EPO standard and the results are set out in graphic form in FIG. 1. Briefly, the results expectedly revealed that the

26

recombinant monkey EPO significantly competed for anti-human EPO antibody although it was not able to completely inhibit binding under the test conditions. The maximum percent inhibition values for recombinant human EPO, however, closely approximated those of the human EPO standard. The parallel nature of the dose response curves suggests immunological identity of the sequences (epitopes) in common. Prior estimates of monkey EPO in culture fluids were re-evaluated at these higher dilution levels and were found to range from 2.91 to 3.12 U/ml. Estimated human EPO production levels were correspondingly set at 392 mU/ml for the five-day growth sample and 567 mU/ml for the seven day growth sample. Estimated monkey EPO production levels in the Example 7B expression system were on the same order or better.

#### EXAMPLE 9

Culture fluids prepared according to Examples 6 and 7 were subjected to an in vitro assay for EPO activity according to the procedure of Goldwasser, et al., *Endocrinology*, 97, 2, pp. 315-323 (1975). Estimated monkey EPO values for culture fluids tested ranged from 3.2 to 4.3 U/ml. Human EPO culture fluids were also active in this in vitro assay and, further, this activity could be neutralized by anti-EPO antibody. The recombinant monkey EPO culture fluids according to Example 6 were also subjected to an assay for in vivo biological activity according to the general procedures of Cotes, et al., *Nature*, 191, pp. 1065-1067 (1961) and Hammond, et al., *Ann.N.Y.Acad.Sci.*, 149, pp. 516-527 (1968) and activity levels ranged from 0.94 to 1.24 U/ml.

#### EXAMPLE 10

In the previous examples, recombinant monkey or human EPO material was produced from vectors used to transfect COS-1 cells. These vectors replicate in COS-1 cells due to the presence of SV40 T antigen within the cell and an SV40 origin of replication on the vectors. Though these vectors produce useful quantities of EPO in COS-1 cells, expression is only transient (7 to 14 days) due to the eventual loss of the vector. Additionally, only a small percentage of COS-1 became productively transfected with the vectors. The present example describes expression systems employing Chinese hamster ovary (CMO) DHFR<sup>-</sup> cells and the selectable marker, DHFR. [For discussion of related expression systems, see U.S. Pat. No. 4,399,216 and European Patent Application Nos. 117058, 117059 and 117060, all published Aug. 29, 1984.]

CHO DHFR<sup>-</sup> cells (DuX-B11) CHO K1 cells, Urlaub, et al., *Proc. Nat. Acad. Sci. (U.S.A.)*, Vol. 77, 4461 (1980) lack the enzyme dihydrofolate reductase (DHFR) due to mutations in the structural genes and therefore require the presence of glycine, hypoxanthine, and thymidine in the culture media. Plasmids pDSVL-MkE (Example 6) or pDSVL-gHuEPO (Example 7B) were transfected along with carrier DNA into CHO DHFR<sup>-</sup> cells growing in media containing hypoxanthine, thymidine, and glycine in 60 mm culture plates. Plasmid pSVgHuEPO (Example 7A) was mixed with the plasmid pMG2 containing a mouse dihydrofolate reductase gene cloned into the bacterial plasmid vector pBR322 (per Gasser, et al., supra). The plasmid mixture and carrier DNA was transfected into CHO DHFR<sup>-</sup> cells. (Cells which acquire one plasmid will generally also acquire a second plasmid). After three days, the cells were dispersed by trypsinization into several 100

4,703,008

27

mm culture plates in media lacking hypoxanthine and thymidine. Only those cells which have been stably transformed with the DHFR gene, and thereby the EPO gene, survive in this media. After 7-21 days, colonies of surviving cells became apparent. These transformant colonies, after dispersion by trypsinization can be continuously propagated in media lacking hypoxanthine and thymidine, creating new cell strains (e.g., CHO pDSVL-MkEPO, CHO pSVgHuEPO, CHO pDSVL-gHuEPO).

Culture fluids from the above cell strains were tested in the RIA for the presence of recombinant monkey or human EPO. Media for strain CHO pDSVL-MkEPO contained EPO with immunological properties like that obtained from COS-1 cells transfected with plasmid pDSVL-MkEPO. A representative 65 hour culture fluid contained monkey EPO at 0.60 U/ml.

Culture fluids from CHO pSVgHuEPO and CHO pDSVL-gHuEPO contained recombinant human EPO with immunological properties like that obtained with COS-1 cells transfected with plasmid pSVgHuEPO or pDSVL-gHuEPO. A representative 3 day culture fluid from CHO pSVgHuEPO contained 2.99 U/ml of human EPO and a 5.5 day sample from CHO pDSVL-gHuEPO had 18.2 U/ml of human EPO as measured by the RIA.

The quantity of EPO produced by the cell strains described above can be increased by gene amplification giving new cell strains of greater productivity. The enzyme dihydrofolate reductase (DHFR) which is the product coded for by the DHFR gene can be inhibited by the drug methotrexate (MTX). More specifically, cells propagated in media lacking hypoxanthine and thymidine are inhibited or killed by MTX. Under the appropriate conditions, (e.g., minimal concentrations of MTX) cells resistant to and able to grow in MTX can be obtained. These cells are found to be resistant to MTX due to an amplification of the number of their DHFR genes, resulting in increased production of DHFR enzyme. The surviving cells can, in turn, be treated with increasing concentrations of MTX, resulting in cell strains containing greater numbers of DHFR genes. "Passenger genes" (e.g., EPO) carried on the expression vector along with the DHFR gene or transformed with the DHFR gene are frequently found also to be increased in their gene copy number.

As examples of practice of this amplification system, cell strain CHO pDSVL-MkE was subjected to increasing MTX concentrations (0 nM, 30 nM and 100 nM). Representative 65-hour culture media samples from each amplification step were assayed by RIA and determined to contain 0.60, 2.45 and 6.10 U/ml, respectively. Cell strain CHO pDSVL-gHuEPO was subjected to a series of increasing MTX concentrations of 30 nM, 50 nM, 100 nM, 200 nM, 1  $\mu$ M, and 5  $\mu$ M MTX. A representative 3-day culture media sample from the 100 nM MTX step contained human EPO at  $3089 \pm 129$  u/ml as judged by RIA. Representative 48 hour cultural medium samples from the 100 nM and 1  $\mu$ M MTX steps contained, respectively, human EPO at 466 and 1352 U/ml as judged by RIA (average of triplicate assays). In these procedures,  $1 \times 10^6$  cells were plated in 5 ml of media in 60 mm culture dishes. Twenty-four hours later the media were removed and replaced with 5 ml of serum-free media (high glucose DMEM supplemented with 0.1 mM non-essential amino acids and L-glutamine). EPO was allowed to accumulate for 48 hours in the serum-free media. The media was collected for RIA

28

assay and the cells were trypsinized and counted. The average RIA values of 467 U/ml and 1352 U/ml for cells grown at 100 nM and 1  $\mu$ M MTX, respectively, provided actual yields of 2335 U/plate and 6750 U/plate. The average cell numbers per plate were  $1.94 \times 10^6$  and  $3.12 \times 10^6$  cells, respectively. The effective production rates for these culture conditions were thus 1264 and 2167 U/ $10^6$  cells/48 hours.

The cells in the cultures described immediately above are a genetically heterogeneous population. Standard screening procedures are being employed in an attempt to isolate genetically homogeneous clones with the highest production capacity. See, Section A, Part 2, of "Points to Consider in the Characterization of Cell Lines Used to Produce Biologics", June 1, 1984, Office of Biologics Research Review, Center for Drugs and Biologics, U.S. Food and Drug Administration.

The productivity of the EPO producing CHO cell lines described above can be improved by appropriate cell culture techniques. The propagation of mammalian cells in culture generally requires the presence of serum in the growth media. A method for production of erythropoietin from CHO cells in media that does not contain serum greatly facilitates the purification of erythropoietin from the culture medium. The method described below is capable of economically producing erythropoietin in serum-free media in large quantities sufficient for production.

Strain CHO pDSVL-gHuEPO cells, grown in standard cell culture conditions, are used to seed spinner cell culture flasks. The cells are propagated as a suspension cell line in the spinner cell culture flask in media consisting of a 50-50 mixture of high glucose DMEM and Ham's F12 supplemented with 5% fetal calf serum, L-glutamine, Penicillin and Streptomycin, 0.05 mM non-essential amino acids and the appropriate concentration of methotrexate. Suspension cell culture allows the EPO-producing CHO cells to be expanded easily to large volumes. CHO cells, grown in suspension, are used to seed roller bottles at an initial seeding density of  $1.5 \times 10^7$  viable cells per 850 cm<sup>2</sup> roller bottle in 200 ml of media. The cells are allowed to grow to confluency as an adherent cell line over a three-day period. The media used for this phase of the growth is the same as used for growth in suspension. At the end of the three-day growth period, the serum containing media is removed and replaced with 100 ml of serum-free media; 50-50 mixture of high glucose DMEM and Ham's F12 supplemented with 0.05 mM non-essential amino acids and L-glutamine. The roller bottles are returned to the roller bottle incubator for a period of 1-3 hours and the media again is removed and replaced with 100 ml of fresh serum-free media. The 1-3 hour incubation of the serum-free media reduces the concentration of contaminating serum proteins. The roller bottles are returned to the incubator for seven days during which erythropoietin accumulates in the serum-free culture media. At the end of the seven-day production phase, the conditioned media is removed and replaced with fresh serum-free medium for a second production cycle. As an example of the practice of this production system, a representative seven-day, serum-free media sample contained human erythropoietin at  $3892 \pm 409$  U/ml as judged by the RIA. Based on an estimated cell density of 0.9 to  $1.8 \times 10^5$  cells/cm<sup>2</sup>, each 850 cm<sup>2</sup> roller bottle contained from 0.75 to  $1.5 \times 10^8$  cells and thus the rate of production of EPO in the 7-day, 100 ml culture was 750 to 1470 U/ $10^6$  cells/48 hours.

4,703,008

29

Culture fluids from cell strain CMO pDSVL-MkEPO carried in 10 nM MTX were subjected to RIA in vitro and in vivo EPO activity assays. The conditioned media sample contained  $41.2 \pm 1.4$  U/ml of MkEPO as measured by the RIA,  $41.2 \pm 0.064$  U/ml as measured by the in vitro biological activity assay and  $42.5 \pm 5$  U/ml as measured by the in vivo biological activity assay. Amino acid sequencing of polypeptide products revealed the presence of EPO products, a principle species having 3 residues of the "leader" sequence adjacent the putative amino terminal alanine. Whether this is the result of incorrect membrane processing of the polypeptide in CHO cells or reflects a difference in structure of the amino terminus of monkey EPO vis-a-vis human EPO, is presently unknown.

Culture fluids from cell strain CHO pDSVL-gHuEPO were subjected to the three assays. A 5.5 day sample contained recombinant human EPO in the media at a level of 18.2 U/ml by RIA assay,  $15.8 \pm 4.6$  U/ml by in vitro assay and  $16.8 \pm 3.0$  U/ml by in vivo assay.

Culture fluid from CHO pDSVL-gHuEPO cells prepared amplified by stepwise 100 nM MTX were subjected to the three assays. A 3.0 day sample contained recombinant human EPO at a level of  $3089 \pm 129$  U/ml by RIA,  $2589 \pm 71.5$  U/ml by in vitro assay, and  $2040 \pm 160$  U/ml by in vivo assay. Amino acid sequencing of this product reveals an amino terminal corresponding to that designated in FIG. 6.

Cell conditioned media from CHO cells transfected with plasmid pDSVL-MkE in 10 nM MTX were pooled, and the MTX dialyzed out over several days, resulting in media with an EPO activity of  $221 \pm 5.1$  U/ml (EPO-CCM). To determine the in vivo effect of the EPO-CCM upon hematocrit levels in normal Balb/C mice, the following experiment was conducted. Cell conditioned media from untransfected CHO cells (CCM) and EPO-CCM were adjusted with PBS. CCM was used for the control group (3 mice) and two dose levels of EPO-CCM—4 units per injection and 44 units per injection—were employed for the experimental groups (2 mice/group). Over the course of 5 weeks, the seven mice were injected intraperitoneally, 3 times per week. After the eighth injection, average hematocrit values for the control group were determined to be 50.4%; for the 4U group, 55.1%; and, for the 44U group, 67.9%.

Mammalian cell expression products may be readily recovered in substantially purified form from culture media using HPLC (C<sub>4</sub>) employing an ethanol gradient, preferably at pH7.

A preliminary attempt was made to characterize recombinant glycoprotein products from conditioned medium of COS-1 and CHO cell expression of the human EPO gene in comparison to human urinary EPO isolates using both Western blot analysis and SDS-PAGE. These studies indicated that the CHO-produced EPO material had a somewhat higher molecular weight than the COS-1 expression product which, in turn, was slightly larger than the pooled source human urinary extract. All products were somewhat heterogeneous. Neuraminidase enzyme treatment to remove sialic acid resulted in COS-1 and CHO recombinant products of approximately equal molecular weight which were both nonetheless larger than the resulting asialo human urinary extract. Endoglycosidase F enzyme (EC 3.2.1) treatment of the recombinant CHO product and the urinary extract product (to totally remove carbohy-

30

drate from both) resulted in substantially homogeneous products having essentially identical molecular weight characteristics.

Purified human urinary EPO and a recombinant, CHO cell-produced, EPO according to the invention were subjected to carbohydrate analysis according to the procedure of Ledeen, et al. *Methods in Enzymology*, 83(Part D), 139-191 (1982) as modified through use of the hydrolysis procedures of Nesser, et al., *Anal. Biochem.*, 142, 58-67 (1984). Experimentally determined carbohydrate constitution values (expressed as molar ratios of carbohydrate in the product) for the urinary isolate were as follows: Hexoses, 1.73; N-acetylglucosamine, 1; N-acetylneuraminic acid, 0.93; Fucose, 0; and N-acetylgalactosamine, 0. Corresponding values for the recombinant product (derived from CHO pDSVL-gHuEPO 3-day culture media at 100 nM MTX) were as follows: Hexoses, 15.09; N-acetylglucosamine, 1; N-acetylneuraminic acid, 0.998; Fucose, 0; and N-acetylgalactosamine, 0. These findings are consistent with the Western blot and SDS-PAGE analysis described above.

Glycoprotein products provided by the present invention are thus comprehensive of products having a primary structural conformation sufficiently duplicative of that of a naturally-occurring erythropoietin to allow possession of one or more of the biological properties thereof and having an average carbohydrate composition which differs from that of naturally-occurring erythropoietin.

#### EXAMPLE 11

The present example relates to the total manufacture by assembly of nucleotide bases of two structural genes encoding the human species EPO sequence of FIG. 6 and incorporating, respectively "preferred" codons for expression in *E. coli* and yeast (*S. cerevisiae*) cells. Also described is the construction of genes encoding analogs of human EPO. Briefly stated, the protocol employed was generally as set out in the previously noted disclosure of Alton, et al. (WO 83/04053). The genes were designed for initial assembly of component oligonucleotides into multiple duplexes which, in turn, were assembled into three discrete sections. These sections were designed for ready amplification and, upon removal from the amplification system, could be assembled sequentially or through a multiple fragment ligation in a suitable expression vector.

FIGS. 10 through 15 and 7 illustrate the design and assembly of a manufactured gene encoding a human EPO translation product lacking any leader or presequence but including an initial methionine residue at position -1. Moreover, the gene incorporated in substantial part *E. coli* preference codons and the construction was therefore referred to as the "ECEPO" gene.

More particularly, FIG. 10 illustrates oligonucleotides employed to generate the Section 1 of the ECEPO gene encoding amino terminal residues of the human species polypeptide. Oligonucleotides were assembled into duplexes (1 and 2, 3 and 4, etc.) and the duplexes were then ligated to provide ECEPO Section 1 as in FIG. 11. Note that the assembled section includes respective terminal EcoRI and BamHI sticky ends, that "downstream" of the EcoRI sticky end is a XbaI restriction enzyme recognition site; and that "upstream" of the BamHI sticky end is a KpnI recognition site. Section 1 could readily be amplified using the M13 phage vector employed for verification of sequence of the section. Some difficulties were encountered in iso-

31

lating the section as an XbaI/KpnI fragment from RF DNA generated in *E. coli*, likely due to methylation of the KpnI recognition site bases within the host. Single-

4,703,008

32

through the second base of the Arg<sup>10</sup> codon. A XbaI/XhoI "linker" sequence was manufactured having the following sequence:

XbaI	+1	2	7	8	9	XhoI
5'-CTAG	Met ATG	Ala GCT	Asn AAT	Cys TGC	Asp GAC-3'	
3'	-TAC	CGA	TTA	ACG	CTG	AGCT-5'

stranded phage DNA was therefore isolated and rendered into double-stranded form in vitro by primer extension and the desired double-stranded fragment was thereafter readily isolated.

ECEPO gene Sections 2 and 3 (FIGS. 13 and 15) were constructed in a similar manner from the oligonucleotides of FIGS. 12 and 14, respectively. Each section was amplified in the M13 vector employed for sequence verification and was isolated from phage DNA. As is apparent from FIG. 13, ECEPO Section 2 was constructed with EcoRI and BamHI sticky ends and could be isolated as a KpnI/BglII fragment. Similarly, ECEPO Section 3 was prepared with BamHI and SalI

The XbaI/XhoI linker and the XhoI/HindIII ECEPO gene sequence fragment were inserted into the large fragment resulting from XbaI and HindIII digestion of plasmid pCFM526—a derivative of plasmid pCFM414 (A.T.C.C. 40076)—as described in co-pending U.S. patent application Ser. No. 636,727, filed Aug. 6, 1984, by Charles F. Morris, to generate a plasmid-borne DNA sequence encoding *E. coli* expression of the Met<sup>-1</sup> form of the desired analog.

#### B. [His<sup>7</sup>]hEPO

Plasmid 536 was digested with HindIII and XhoI as in part A above. A XbaI/XhoI linker was manufactured having the following sequence:

XbaI	+1	2	3	4	5	6	7	8	9	XhoI
5'-CTAG	Met ATG	Ala GCT	Pro CCG	Pro CCA	Arg CGT	Leu CTG	Ile ATC	His CAT	Asp GAC-3'	
3'	-TAC	CGA	GGC	GGT	GCA	GAC	TAG	GTA	CTG	AGCT-5'

sticky ends and could be isolated from phage RF DNA as a BglII/SalI fragment. The three sections thus prepared can readily be assembled into a continuous DNA sequence (FIG. 7) encoding the entire human species EPO polypeptide with an amino terminal methionine codon (ATG) for *E. coli* translation initiation. Note also that "upstream" of the initial ATG is a series of base pairs substantially duplicating the ribosome binding site sequence of the highly expressed OMP-f gene of *E. coli*.

Any suitable expression vector may be employed to carry the ECEPO. The particular vector chosen for expression of the ECEPO gene as the "temperature sensitive" plasmid pCFM536—a derivative of plasmid pCFM414 (A.T.C.C. 40076)—as described in co-pending U.S. patent application Ser. No. 636,727, filed Aug. 6, 1984, (published EPO Application No. 136,490) by Charles F. Morris. More specifically, pCFM536 was digested with XbaI and HindIII; the large fragment was isolated and employed in a two-part ligation with the ECEPO gene. Sections 1 (XbaI/KpnI), 2 (KpnI/BglII) and 3 (BglII/SalI) had previously been assembled in the correct order in M13 and the EPO gene was isolated therefrom as a single XbaI/HindIII fragment. This fragment included a portion of the polylinker from M13 mp9 phage spanning the SalI to HindIII sites therein. Control of expression in the resulting expression plasmid, p536, was by means of a lambda P<sub>L</sub> promoter, which itself may be under control of the C<sub>J857</sub> repressor gene (such as provided in *E. coli* strain K12ΔHtrp).

The manufactured ECEPO gene above may be variously modified to encode erythropoietin analogs such as [Asn<sup>2</sup>, des-Pro<sup>2</sup> through Ile<sup>6</sup>]hEPO and [His<sup>7</sup>]hEPO, as described below.

#### A. [Asn<sup>2</sup>, des-Pro<sup>2</sup> through Ile<sup>6</sup>]hEPO

Plasmid 536 carrying the ECEPO manufactured gene of FIG. 7 as a XbaI to HindIII insert was digested with HindIII and XhoI. The latter endonuclease cuts the ECEPO gene at a unique, 6 base pair recognition site spanning the last base of the codon encoding Asp<sup>8</sup>

The linker and the XhoI/HindIII ECEPO sequence fragment were then inserted into pCFM526 to generate a plasmid-borne DNA sequence encoding *E. coli* expression of the Met<sup>-1</sup> form of the desired analog.

Construction of a manufactured gene ("SCEPO") incorporating yeast preference codons is as described in the following FIGS. 16 through 21 and 8. As was the case with the ECEPO gene, the entire construction involved formation of three sets of oligonucleotides (FIGS. 16, 18 and 20) which were formed into duplexes and assembled into sections (FIGS. 17, 19 and 21). Note that synthesis was facilitated in part by use of some sub-optimal codons in both the SCEPO and ECEPO constructions, i.e., oligonucleotides 7–12 of Section 1 of both genes were identical, as were oligonucleotides 1–6 of Section 2 in each gene.

The assembled SCEPO sections were sequenced in M13 and Sections 1, 2 and 3 were isolatable from the phage as HindIII/KpnI, KpnI/BglII, and BglII/SalI fragments.

The presently preferred expression system for SCEPO gene products is a secretion system based on *S. cerevisiae*  $\alpha$ -factor secretion, as described in co-pending U.S. patent application Ser. No. 487,753, filed Apr. 22, 1983, by Grant A. Bitter, published Oct. 31, 1984 as European Patent Application No. 0 123 294. Briefly put, the system involves constructions wherein DNA encoding the leader sequence of the yeast  $\alpha$ -factor gene product is positioned immediately 5' to the coding region of the exogenous gene to be expressed. As a result, the gene product translated includes a leader or signal sequence which is "processed off" by an endogenous yeast enzyme in the course of secretion of the remainder of the product. Because the construction makes use of the  $\alpha$ -factor translation initiation (ATG) codon, there was no need to provide such a codon at the -1 position of the SCEPO gene. As may be noted from FIG. 8, the

4,703,008

33

alanine (+1) encoding sequence is preceded by a linker sequence allowing for direct insertion into a plasmid including the DNA for the first 80 residues of the  $\alpha$ -factor leader following the  $\alpha$ -factor promoter. The specific preferred construction for SCEPO gene expression involved a four-part ligation including the above-noted SCEPO section fragments and the large fragment of HindIII/SalI digestion of plasmid p $\alpha$ C3. From the resulting plasmid p $\alpha$ C3/SCEPO, the  $\alpha$ -factor promoter and leader sequence and SCEPO gene were isolated by digestion with BamHI and ligated into BamHI digested plasmid pYE to form expression plasmid pYE/SCEPO.

## EXAMPLE 12

The present example relates to expression of recombinant products of the manufactured ECEPO and SCEPO genes within the expression systems of Example 11.

In use of the expression system designed for use of *E. coli* host cells, plasmid p536 of Example 11 was transformed into AM7 *E. coli* cells previously transformed with a suitable plasmid, pMW1, harboring a  $C_{\beta 857}$  gene. Cultures of cells in LB broth (Ampicillin 50  $\mu$ g/ml and kanamycin 5  $\mu$ g/ml, preferably with 10 mM  $MgSO_4$ ) were maintained at 28° C. and upon growth of cells in culture to O.D.<sub>600</sub>=0.1, EPO expression was induced by raising the culture temperature to 42° C. Cells grown to about 40 O.D. provided EPO production (as estimated by gel) of about 5 mg/OD liter.

Cells were harvested, lysed, broken with French Press (10,000 psi) and treated with lysozyme and NP-40 detergent. The pellet resulting from 24,000 xg centrifugation was solubilized with guanidine HCl and subjected to further purification in a single step by means of C<sub>4</sub> (Vydac) Reverse Phase HPLC (EtOH, 0-80%, 50 mM  $NH_4Ac$ , pH 4.5). Protein sequencing revealed the product to be greater than 95% pure and the products obtained revealed two different amino terminals, A-P-P-R . . . and P-P-R . . . in a relative quantitative ratio of about 3 to 1. This latter observation of hEPO and [des Ala<sup>1</sup>]hEPO products indicates that amino terminal "processing" within the host cells serves to remove the terminal methionine and in some instances the initial alanine. Radioimmunoassay activity for the isolates was at a level of 150,000 to 160,000 U/mg; in vitro assay activity was at a level of 30,000 to 62,000 U/mg; and in vivo assay activity ranged from about 120 to 720 U/mg. (Cf., human urinary isolate standard of 70,000 U/mg in each assay.) The dose response curve for the recombinant product in the in vivo assay differed markedly from that of the human urinary EPG standard.

The EPO analog plasmids formed in parts A and B of Example 11 were each transformed into pMW1-transformed AM7 *E. coli* cells and the cells were cultured as above. Purified isolates were tested in both RIA and in vitro assays. RIA and in vitro assay values for [Asn<sup>2</sup>, des-Pro<sup>2</sup> through Ile<sup>6</sup>]hEPO expression products were approximately 11,000 U/mg and 6,000 U/mg protein, respectively, while the assay values for [His<sup>7</sup>]hEPO were about 41,000 U/mg and 14,000 U/mg protein, respectively, indicating that the analog products were from one-fourth to one-tenth as "active" as the "parent" expression product in the assays.

In the expression system designed for use of *S. cerevisiae* host cells, plasmid pYE/SCEPO was transformed into two different strains, YSDP4 (genotype  $\alpha$  pep4-3 trpl) and RK81 (genotype  $\alpha$ apep4-3 trpl). Transformed YSDP4 hosts were grown in SD medium (Methods in

34

Yeast Genetics, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., p. 62 (1983) supplemented with casamino acids at 0.5%, pH 6.5 at 30° C. Media harvested when the cells had been grown to 36 O.D. contained EPO products at levels of about 244 U/ml (97  $\mu$ g/OD liter by RIA). Transformed RK81 cells grown to either 6.5 O.D. or 60 O.D. provided media with EPO concentrations of about 80-90 U/ml (34  $\mu$ g/OD liter by RIA). Preliminary analyses reveal significant heterogeneity in products produced by the expression system, likely to be due to variations in glycosylation of proteins expressed, and relatively high mannose content of the associated carbohydrate.

Plasmids p $\alpha$ C3 and pYE in HB101 *E. coli* cells were deposited in accordance with the Rules of Practice of the U.S. Patent Office on Sept. 27, 1984, with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Md., under deposit numbers A.T.C.C. 39881 and A.T.C.C. 39882, respectively. Plasmids pCFM526 in AM7 cells, pCFM536 in JM103 cells, and pMW1 in JM103 cells were likewise deposited on Nov. 21, 1984 as A.T.C.C. 33932, 33934, and 33933, respectively. *Saccharomyces cerevisiae* strains YSDP4 and RK81 were deposited on Nov. 21, 1984 as A.T.C.C. 20734 and 20733, respectively.

It should be readily apparent from consideration of the above illustrative examples that numerous exceptionally valuable products and processes are provided by the present invention in its many aspects.

Polypeptides provided by the invention are conspicuously useful materials, whether they are microbially expressed products or synthetic products, the primary, secondary or tertiary structural conformation of which was first made known by the present invention.

As previously indicated, recombinant-produced and synthetic products of the invention share, to varying degrees, the in vitro biological activity of EPO isolates from natural sources and consequently are projected to have utility as substitutes for EPO isolates in culture media employed for growth of erythropoietic cells in culture. Similarly, to the extent that polypeptide products of the invention share the in vivo activity of natural EPO isolates they are conspicuously suitable for use in erythropoietin therapy procedures practiced on mammals, including humans, to develop any or all of the effects herefore attributed in vivo to EPO, e.g., stimulation of reticulocyte response, development of ferrokinetic effects (such as plasma iron turnover effects and marrow transit time effects), erythrocyte mass changes, stimulation of hemoglobin C synthesis (see, Eschbach, et al., supra) and, as indicated in Example 10, increasing hematocrit levels in mammals. Included within the class of humans treatable with products of the invention are patients generally requiring blood transfusions and including trauma victims, surgical patients, renal disease patients including dialysis patients, and patients with a variety of blood composition affecting disorders, such as hemophilia, sickle cell disease, physiologic anemias, and the like. The minimization of the need for transfusion therapy through use of EPO therapy can be expected to result in reduced transmission of infectious agents. Products of the invention, by virtue of their production by recombinant methods, are expected to be free of pyrogens, natural inhibitory substances, and the like, and are thus likely to provide enhanced overall effectiveness in therapeutic processes vis-a-vis naturally derived products. Erythropoietin therapy with products of the present invention is also expected to be use-



4,703,008

35

ful in the enhancement of oxygen carrying capacity of individuals encountering hypoxic environmental conditions and possibly in providing beneficial cardiovascular effects.

A preferred method for administration of polypeptide products of the invention is by parenteral (e.g., IV, IM, SC, or IP) routes and the compositions administered would ordinarily include therapeutically effective amounts of product in combination with acceptable diluents, carriers and/or adjuvants. Preliminary pharmacokinetic studies indicate a longer half-life in vivo for monkey EPO products when administered IM rather than IV. Effective dosages are expected to vary substantially depending upon the condition treated but therapeutic doses are presently expected to be in the range of 0.1 (~7U) to 100 (~7000U)  $\mu\text{g}/\text{kg}$  body weight of the active material. Standard diluents such as human serum albumin are contemplated for pharmaceutical compositions of the invention, as are standard carriers such as saline.

Adjuvant materials suitable for use in compositions of the invention include compounds independently noted for erythropoietic stimulatory effects, such as testosterone, progenitor cell stimulators, insulin-like growth factor, prostaglandins, serotonin, cyclic AMP, prolactin and triiodothyronine, as well as agents generally employed in treatment of aplastic anemia, such as methenolene, stanozolol and nandrolone [see, e.g., Resegotti, et al., *Panminerva Medica*, 23, 243-248 (1981); McGonigle, et al., *Kidney Int.*, 25(2), 437-444 (1984); Pavlovic-Kantera, et al., *Expt. Hematol.*, 8(Supp. 8), 283-291 (1980); and Kurtz, *FEBS Letters*, 14a(1), 105-108 (1982)]. Also contemplated as adjuvants are substances reported to enhance the effects of, or synergize, erythropoietin or asialo-EPO, such as the adrenergic agonists, thyroid hormones, androgens and BPA [see, Dunn, "Current Concepts in Erythropoiesis", John Wiley and Sons (Chichester, England, 1983); Weiland, et al., *Blut*, 44(3), 173-175 (1982); Kalmanti, *Kidney Int.*, 22, 383-391 (1982); Shahidi, *New Eng. J. Med.*, 289, 72-80 (1973); Fisher, et al., *Steroids*, 30(6), 833-845 (1977); Urabe, et al., *J. Exp. Med.*, 149, 1314-1325 (1979); and Billat, et al., *Expt. Hematol.*, 10(1), 133-140 (1982)] as well as the classes of compounds designated "hepatic erythropoietic factors" [see, Naughton, et al., *Acta Haemat.*, 69, 171-179 (1983)] and "erythropoietins" [as described by Congote, et al. in Abstract 364, Proceedings 7th International Congress of Endocrinology (Quebec City, Quebec, July 1-7, 1984); Congote, *Biochem. Biophys. Res. Comm.*, 115(2), 447-483 (1983) and Congote, *Anal. Biochem.*, 140, 428-433 (1984)] and "erythroge-nins" [as described in Rothman, et al., *J. Surg. Oncol.*, 20, 105-108 (1982)]. Preliminary screenings designed to measure erythropoietic responses of ex-hypoxic polycythemic mice pre-treated with either 5- $\alpha$ -dihydrotestosterone or nandrolone and then given erythropoietin of the present invention have generated equivocal results.

Diagnostic uses of polypeptides of the invention are similarly extensive and include use in labelled and unlabelled forms in a variety of immunoassay techniques including RIA's, ELISA's and the like, as well as a variety of in vitro and in vivo activity assays. See, e.g., Dunn, et al., *Expt. Hematol.*, 11(7), 590-600 (1983); Gibson, et al., *Pathology*, 16, 155-156 (1984); Krystal, *Expt. Hematol.*, 11(7), 649-660 (1983); Saito, et al., *Jap. J. Med.*, 23(1), 16-21 (1984); Nathan, et al., *New Eng. J. Med.*, 308(9), 520-522 (1983); and various references pertaining to assays referred to therein. Polypep-

36

tides of the invention, including synthetic peptides comprising sequences of residues of EPO first revealed herein, also provide highly useful pure materials for generating polyclonal antibodies and "banks" of monoclonal antibodies specific for differing continuous and discontinuous epitopes of EPO. As one example, preliminary analysis of the amino acid sequences of FIG. 6 in the context of hydropathicity according to Hopp, et al., *P.N.A.S. (U.S.A.)*, 78, pp. 3824-3828 (1981) and of secondary structures according to Chou, et al., *Ann. Rev. Biochem.*, 47, p. 251 (1978) revealed that synthetic peptides duplicative of continuous sequences of residues spanning positions 41-57 inclusive, 116-128 inclusive and 144-166 inclusive are likely to produce a highly antigenic response and generate useful monoclonal and polyclonal antibodies immunoreactive with both the synthetic peptide and the entire protein. Such antibodies are expected to be useful in the detection and affinity purification of EPO and EPO-related products.

Illustratively, the following three synthetic peptides were prepared:

- (1) hEPO 41-57, V-P-D-T-K-V-N-F-Y-A-W-K-R-M-E-V-G;
- (2) hEPO 116-128, K-E-A-I-S-P-P-D-A-A-S-A-A;
- (3) hEPO 144-166, V-Y-S-N-F-L-R-G-K-L-K-L-Y-T-G-E-A-C-R-T-G-D-R.

Preliminary immunization studies employing the above-noted polypeptides have revealed a relatively weak positive response to hEPO 41-57, no appreciable response to hEPO 116-128, and a string positive response to hEPO 144-166, as measured by capacity of rabbit serum antibodies to immunoprecipitate <sup>125</sup>I-labelled human urinary EPO isolates. Preliminary in vivo activity studies on the three peptides revealed no significant activity either alone or in combination.

While the deduced sequences of amino acid residues of mammalian EPO provided by the illustrative examples essentially define the primary structural conformation of mature EPO, it will be understood that the specific sequence of 165 amino acid residues of monkey species EPO in Table V and the 166 residues of human species EPO in FIG. 6 do not limit the scope of useful polypeptides provided by the invention. Comprehended by the present invention are those various naturally-occurring allelic forms of EPO which past research into biologically active mammalian polypeptides such as human  $\gamma$  interferon indicates are likely to exist. (Compare, e.g., the human immune interferon species reported to have an arginine residue at position No. 140 in EPO published application No. 0 077 670 and the species reported to have glutamine at position No. 140 in Gray, et al., *Nature*, 295, pp. 503-508 (1982). Both species are characterized as constituting "mature" human  $\gamma$  interferon sequences.) Allelic forms of mature EPO polypeptides may vary from each other and from the sequences of FIG. 5 and 6 in terms of length of sequence and/or in terms of deletions, substitutions, insertions or additions of amino acids in the sequence, with consequent potential variations in the capacity for glycosylation. As noted previously, one putative allelic form of human species EPO is believed to include a methionine residue at position 126. Expectedly, naturally-occurring allelic forms of EPO-encoding DNA genomic and cDNA sequences are also likely to occur which code for the above-noted types of allelic polypeptides or simply employ differing codons for designation of the same polypeptides as specified.

4,703,008

37

In addition to naturally-occurring allelic forms of mature EPO, the present invention also embraces other "EPO products" such as polypeptide analogs of EPO and fragments of "mature" EPO. Following the procedures of the above-noted published application by Alton, et al. (WO/83/04053) one may readily design and manufacture genes coding for microbial expression of polypeptides having primary conformations which differ from that herein specified for mature EPO in terms of the identity or location of one or more residues (e.g., substitutions, terminal and intermediate additions and deletions). Alternately, modifications of cDNA and genomic EPO genes may be readily accomplished by well-known site-directed mutagenesis techniques and employed to generate analogs and derivatives of EPO. Such EPO products would share at least one of the biological properties of EPO but may differ in others. As examples, projected EPO products of the invention include those which are foreshortened by e.g., deletions 8 Asn<sup>2</sup>, des-Pro<sup>2</sup> through Ile<sup>6</sup>hEPO, [des-Thr<sup>163</sup> through Arg<sup>166</sup>]hEPO and "Δ27-55hEPO", the latter having the residues coded for by an entire exon deleted; or which are more stable to hydrolysis (and, therefore, may have more pronounced or longer lasting effects than naturally-occurring EPO); or which have been altered to delete one or more a potential sites for glycosylation (which may result in higher activities for yeast-produced products); or which have one or more cysteine residues deleted or replaced by, e.g., histidine or serine residues (such as the analog [His<sup>7</sup>]hEPO) and are potentially more easily isolated in active form from microbial systems; or which have one or more tyrosine residues replaced by phenylalanine (such as the analogs [Phe<sup>15</sup>]hEPO, [Phe<sup>49</sup>]hEPO, and [Phe<sup>145</sup>]hEPO) and may bind more or less readily to EPO receptors on target cells. Also comprehended are polypeptide fragments duplicating only a part of the continuous amino acid sequence or secondary conformations within mature EPO, which fragments may possess one activity of EPO (e.g., receptor binding) and not others (e.g., erythropoietic activity). Especially significant in this regard are those potential fragments of EPO which are elucidated upon consideration of the human genomic DNA sequence of Table VI, i.e., "fragments" of the total continuous EPO sequence which are delineated by intron sequences and which may constitute distinct "domains" of biological activity. It is noteworthy that the absence of in vivo activity for any one or more of the "EPO products" of the invention is not wholly preclusive of therapeutic utility (see, Weiland, et al., supra) or of utility in other contexts, such as in EPO assays or EPO antagonism. Antagonists of erythropoietin may be quite useful in treatment of polycythemia or cases of overproduction of EPO [see, e.g., Adamson, *Hosp. Practice*, 18(12), 49-57 (1983), and Hellmann, et al., *Clin. Lab. Haemat.*, 5, 335-342 (1983)].

According to another aspect of the present invention, the cloned DNA sequences described herein which encode human and monkey EPO polypeptides are conspicuously valuable for the information which they provide concerning the amino acid sequence of mammalian erythropoietin which has theretofore been unavailable despite decades of analytical processing of isolates of naturally-occurring products. The DNA sequences are also conspicuously valuable as products useful in effecting the large scale microbial synthesis of erythropoietin by a variety of recombinant techniques. Put another way, DNA sequences provided by the

38

invention are useful in generating new and useful viral and circular plasmid DNA vectors, new and useful transformed and transfected microbial prokaryotic and eucaryotic host cells (including bacterial and yeast cells and mammalian cells grown in culture), and new and useful methods for cultured growth of such microbial host cells capable of expression of EPO and EPO products. DNA sequences of the invention are also conspicuously suitable materials for use as labelled probes in isolating EPO and related protein encoding cDNA and genomic DNA sequences of mammalian species other than human and monkey species herein specifically illustrated. The extent to which DNA sequences of the invention will have use in various alternative methods of protein synthesis (e.g., in insect cells) or in genetic therapy in humans and other mammals cannot yet be calculated. DNA sequences of the invention are expected to be useful in developing transgenic mammalian species which may serve as eucaryotic "hosts" for production of erythropoietin and erythropoietin products in quantity. See, generally, Palmiter, et al., *Science*, 222(4625), 809-814 (1983).

Viewed in this light, therefore, the specific disclosures of the illustrative examples are clearly not intended to be limiting upon the scope of the present invention and numerous modifications and variations are expected to occur to those skilled in the art. As one example, while DNA sequences provided by the illustrative examples include cDNA and genomic DNA sequences, because this application provides amino acid sequence information essential to manufacture of DNA sequence, the invention also comprehends such manufactured DNA sequences as may be constructed based on knowledge of EPO amino acid sequences. These may code for EPO (as in Example 12) as well as for EPO fragments and EPO polypeptide analogs (i.e., "EPO Products") which may share one or more biological properties of naturally-occurring EPO but not share others (or possess others to different degrees).

DNA sequences provided by the present invention are thus seen to comprehend all DNA sequences suitable for use in securing expression in a prokaryotic or eucaryotic host cell of a polypeptide product having at least a part of the primary structural conformation and one or more of the biological properties of erythropoietin, and selected from among: (a) the DNA sequences set out in FIGS. 5 and 6; (b) DNA sequences which hybridize to the DNA sequences defined in (a) or fragments thereof; and (c) DNA sequences which, but for the degeneracy of the genetic code, would hybridize to the DNA sequences defined in (a) and (b). It is noteworthy in this regard, for example, that existing allelic monkey and human EPO gene sequences and other mammalian species gene sequences are expected to hybridize to the sequences of FIGS. 5 and 6 or to fragments thereof. Further, but for the degeneracy of the genetic code, the SCEPO and ECEPO genes and the manufactured or mutagenized cDNA or genomic DNA sequences encoding various EPO fragments and analogs would also hybridize to the above-mentioned DNA sequences. Such hybridizations could really be carried out under the hybridization conditions described herein with respect to the initial isolation of the monkey and human EPO-encoding DNA or more stringent conditions, if desired to reduce background hybridization.

In a like manner, while the above examples illustrate the invention of microbial expression of EPO products

in the context of mammalian cell expression of DNA inserted in a hybrid vector of bacterial plasmid and viral genomic origins, a wide variety of expression systems are within the contemplation of the invention. Conspicuously comprehended are expression systems involving vectors of homogeneous origins applied to a variety of bacterial, yeast and mammalian cells in culture as well as to expression systems not involving vectors (such as calcium phosphate transfection of cells). In this regard, it will be understood that expression of, e.g., monkey origin DNA in monkey host cells in culture and human host cells in culture, actually constitute instances of "exogenous" DNA expression inasmuch as the EPO DNA whose high level expression is sought would not have its origins in the genome of the host. Expression systems of the invention further contemplate these practices resulting in cytoplasmic formation of EPO products and accumulation of glycosylated and non-glycosylated EPO products in host cell cytoplasm or membranes (e.g., accumulation in bacterial periplasmic spaces) or in culture medium supernatants as above illustrated, or in rather uncommon systems such as *P. aeruginosa* expression systems (described in Gray, et al., *Biotechnology*, 2, pp. 161-165 (1984)).

Improved hybridization methodologies of the invention, while illustratively applied above to DNA/DNA hybridization screenings are equally applicable to RNA/RNA and RNA/DNA screening. Mixed probe techniques as herein illustrated generally constitute a number of improvements in hybridization processes allowing for more rapid and reliable polynucleotide isolations. These many individual processing improvements include: improved colony transfer and maintenance procedures; use of nylon-based filters such as GeneScreen and GeneScreen Plus to allow reprobation with same filters and repeated use of the filter, application of novel protease treatments. [compared, e.g., to Taub, et al. *Anal. Biochem.*, 126, pp. 222-230 (1982)]; use of very low individual concentrations (on the order of 0.025 picomole) of a large number of mixed probes (e.g., numbers in excess of 32); and, performing hybridization and post-hybridization steps under stringent temperatures closely approaching (i.e., within 4° C. and preferably within 2° C. away from) the lowest calculated dissociation temperature of any of the mixed probes employed. These improvements combine to provide results which could not be expected to attend their use. This is amply illustrated by the fact that mixed probe procedures involving 4 times the number of probes ever before reported to have been successfully used in even cDNA screens on messenger RNA species of relatively low abundance were successfully applied to the isolation of a unique sequence gene in a genomic library screening of 1,500,000 phage plaques. This feat was accomplished essentially concurrently with the publication of the considered opinion of Anderson, et al., supra, that mixed probe screening methods were "... impractical for isolation of mammalian protein genes when corresponding RNA's are unavailable.

What is claimed is:

1. A purified and isolated DNA sequence encoding erythropoietin, said DNA sequence selected from the group consisting of:

- (a) the DNA sequences set out in FIGS. 5 and 6 or their complementary strands; and
- (b) DNA sequences which hybridize under stringent conditions to the DNA sequences defined in (a).

2. A purified and isolated DNA sequence consisting essentially of a DNA sequence encoding human erythropoietin.

3. A purified and isolated DNA sequence consisting essentially of a DNA sequence encoding monkey erythropoietin.

4. A procaryotic or eucaryotic host cell transformed or transfected with a DNA sequence according to claim 1, 2 or 3 in a manner allowing the host cell to express erythropoietin.

5. A biologically functional circular plasmid or viral DNA vector including a DNA sequence according to claim 1, 2, or 3.

6. A procaryotic or eucaryotic host cell stably transformed or transfected with a DNA vector according to claim 5.

7. A purified and isolated DNA sequence consisting essentially of a DNA sequence encoding a polypeptide having an amino acid sequence sufficiently duplicative of that of erythropoietin to allow possession of the biological property of causing bone marrow cells to increase production of reticulocytes and red blood cells, and to increase hemoglobin synthesis or iron uptake.

8. A cDNA sequence according to claim 7.

9. A monkey species erythropoietin coding DNA sequence according to claim 8.

10. A DNA sequence according to claim 9 and including the protein coding region set forth in FIG. 5.

11. A genomic DNA sequence according to claim 7.

12. A human species erythropoietin coding DNA sequence according to claim 11.

13. A DNA sequence according to claim 12 and including the protein coding region set forth in FIG. 6.

14. A DNA sequence according to claim 7 and including one or more codons preferred for expression in *E. coli* cells.

15. A DNA sequence according to claim 14, coding for expression of human species erythropoietin.

16. A DNA sequence according to claim 15 including the protein coding region set forth in FIG. 7.

17. A DNA sequence according to claim 7 and including one or more codons preferred for expression in yeast cells.

18. A DNA sequence according to claim 17, coding for expression of human species erythropoietin.

19. A DNA sequence according to claim 18 including the protein coding region set forth in FIG. 8.

20. A DNA sequence according to claim 7 covalently associated with a detectable label substance.

21. A DNA sequence according to claim 20 wherein the detectable label is a radiolabel.

22. A single-strand DNA sequence according to claim 20.

23. A procaryotic or eucaryotic host cell transformed or transfected with a DNA sequence according to claim 7, 8, or 11 in a manner allowing the host cell to express said polypeptide.

24. A transformed or transfected host cell according to claim 23 which host cell is capable of glycosylating said polypeptide.

25. A transformed or transfected mammalian host cell according to claim 24.

26. A transformed or transfected COS cell according to claim 25.

27. A transformed or transfected CHO cell according to claim 25.

4,703,008

41

28. A biologically functional circular plasmid or viral DNA vector including a DNA sequence according to claim 7.

29. A procaryotic or eucaryotic host cell stably transformed or transfected with a DNA vector according to claim 28.

30. A DNA sequence according to claim 7 coding for

42

[Phe<sup>15</sup>]hEPO, [Phe<sup>49</sup>]hEPO, [Phe<sup>145</sup>]hEPO, [His<sup>7</sup>]hEPO, [Asn<sup>2</sup>des-Pro<sup>2</sup> through Ile<sup>6</sup>]hEPO, [des-Thr<sup>163</sup> through Arg<sup>166</sup>]hEPO, or [ $\Delta$ 27-55]hEPO.

31. A purified and isolated DNA sequence as set out in FIGS. 5 or 6 or the complementary strand of such a sequence.

\* \* \* \* \*

10

15

20

25

30

35

40

45

50

55

60

65

UNITED STATES PATENT

CERTIFICATE OF CORRECTION

PATENT NO. : 4,703,008

DATED : October 27, 1987

Page 1 of 8

INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Abstract, line 22, "heterologus" should be --heterologous--

Col. 1, line 17, "erythropoletin" should be  
--erythropoietin--

Col. 1, line 33, "nucleotlde" should be --nucleotide--

Col. 1, line 55, "smallRNA" should be --small RNA--

Col. 1, line 62, "grouplngs" should be --groupings--

Col. 1, line 67, "promoter" should be --Promoter--

Col. 2, line 6, "sequenoos" should be --sequences--

Col. 2, line 36, "amolification" should be --amplification--

Col. 2, line 37, please insert "the" after "in"

Col. 2, line 41, "whioh" should be --which--

Col. 2, line 46, please insert ")" after "heterologous"

Col. 2, line 51, "restoratlon" should be --restoration--

Col. 2, line 60, "frequentiy" should be --frequently--

Col. 3, line 24, "WO83/0405" should be --WO83/04053--

Col. 3, line 31, "sequenoos" should be --sequences--

UNITED STATES PATENT AND TRADEMARK OFFICE  
CERTIFICATE OF CORRECTION

PATENT NO. : 4,703,008

DATED : October 27, 1987

Page 2 of 8

INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Col. 3, line 59, "singlestranded" should be  
--single-stranded--

Col. 3, line 63, "tc" should be --to--

Col. 3, line 64, "provlded" should be --provided--

Col. 4, line 12, "techniques:" should be --techniques;--

Col. 4, line 43, "32 member" should be --32-member--

Col. 4, line 47, "DMA" should be --DNA--

Col. 4, line 57, "DMA" should be --DNA--

Col. 5, line 4, please delete the second occurrence of "the"

Col. 5, line 9, "80pp." should be --80, pp.--

Col. 5, line 14, "panoreatic" should be --pancreatic--

Col. 5, line 32, "librales" should be --libraries--

Col. 5, line 51, "polypeptide" should be --Polypeptide--

Col. 5, line 64, please insert "but" after "components"

Col. 5, lines 66-67, "Carbohydrate" should be --carbohydrate--

Col. 6, lines 2-3, "Tbis normai" should be --This normal--

UNITED STATES PATENT AND TRADEMARK OFFICE  
CERTIFICATE OF CORRECTION

PATENT NO. : 4,703,008

DATED : October 27, 1987

Page 3 of 8

INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Col. 6, line 23, "Exp.Mematol." should be --Exp.Hematol.--

Col. 6, line 27, "Expt.Hematol." should be --Exp.Hematol.--

Col. 6, line 28, "1980:" should be --(1980);--

Col. 6, line 30, please insert a space before "1832"

Col. 6, line 33, "Desspyris" should be --Dessypris--

Col. 6, line 50, "alo" should be --also--

Col. 7, line 6, please delete the character "I" between  
"1106" and "(1983)"

Col. 7, line 10, "erythropletin" should be  
--erythropoietin--

Col. 7, line 25, "urin" should be --urine--

Col. 7, line 40, please insert a quotation mark (") after  
"effects"

Col. 7, line 47, please insert a close parenthesis ")"  
after "propagation"

Col. 7, line 65, "erythlopoietin" should be --erythropoietin--

Col. 8, line 43, "moiecular" should be --molecular--

Col. 9, lines 14-15, "erythlopoietin" should be  
--erythropoietin--

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 4,703,008

DATED : October 27, 1987

Page 4 of 8

INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

- Col. 10, line 19, "Whlle" should be --While--
- Col. 10, line 50, "characterired" should be --characterized--
- Col. 10, line 64, "mammallan" should be --mammalian--
- Col. 11, line 2, "prlmary" should be --primary--
- Col. 11, line 11, "whioh" should be --which--
- Col. 11, line 46, "polypep-tides" should be --polypeptides--
- Col. 12, line 10, "analoqs" should be --analogs--
- Col. 12, line 17, "ccnformation" should be --conformation--
- Col. 12, line 18, "propezties" should be --properties--
- Col. 12, line 26, "DMA" should be --DNA--
- Col. 12, line 20, "Tables V and VI" should be  
--Figures 5 and 6--
- Col. 12, line 39, "Table VI" should be --Figure 6--
- Col. 12, line 63, "DMA" should be --DNA--
- Col. 12, line 64, "neighboring" should be --neighboring--
- Col. 13, line 67, "construction" should be --constructions--



UNITED STATES PATENT AND TRADEMARK OFFICE  
CERTIFICATE OF CORRECTION

PATENT NO. : 4,703,008

DATED : October 27, 1987

Page 5 of 8

INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Col. 14, line 12, "biological" should be --biological--

Col. 14, line 45, "homology" should be --homology--

Col. 14, line 47, "fragments" should be --fragments--

Col. 14, line 62, "immunological" should be --immunological--

Col. 14, line 64, "lis" should be --is--

Col. 14, line 65, "fragments" should be --fragments--

Col. 15, lines 29-30, "Example" should be --Examples--

Col. 15, line 35, "genomlc" should be --genomic--

Col. 15, line 36, "CMO" should be --CHO--

Col. 15, line 38, "charactezization" should be  
--characterization--

Col. 16, line 43, please insert a parenthesis "(" before "Gln"

Col. 16, line 56, "qamma" should be --gamma--

Col. 17, line 15, please insert "of" before "either"

Col. 17, line 52, please delete the comma after "Springs"

Col. 21, line 25, "(designated  $\lambda_{HE1}$ )" should be  
--(designated  $\lambda_{HE1}$ )--

UNITED STATES PATENT AND TRADEMARK OFFICE  
CERTIFICATE OF CORRECTION

PATENT NO. : 4,703,008

DATED : October 27, 1987

Page 6 of 8

INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Col. 21, line 27, "The" should be --the--

Col. 21, line 41, "glutamine" should be --glutamic acid--

Col. 21, lines 56-57, "Table VI" should be --Figure 6--

Col. 22, line 36, "l.e.," should be --i.e.,--

Col. 22, line 49, "Iinker" should be --linker--

Col. 22, line 52, "BamHMI" should be --BamHI--

Col. 23, line 34, "(DMFR)" should be --(DHFR)--

Col. 23, line 43, "litaged" should be --ligated--

Col. 24, line 39, "EcoRl" should be --EcoRI--

Col. 24, line 52, "BamHl" should be --BamHI--

Col. 24, line 55, "BamHl" should be --BamHI--

Col. 24, line 60, "angalysis" should be --analysis--

Col. 25, line 23, "approxImately" should be --approximately--

Col. 25, line 34, "44" should be --53--

Col. 26, line 46, "(CMO)" should be --(CHO)--

Col. 27, line 24, "snd" should be --and--

Col. 27, line 42, "DMFR" should be --DHFR--

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 4,703,008

DATED : October 27, 1987

Page 7 of 8

INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

- Col. 27, line 54, "serles" should be --series--
- Col. 28, line 12, "hemogeneous" should be --homogeneous--
- Col. 29, line 1, "CMO" should be --CHO--
- Col. 30, line 54, please insert a period "." after "gene"
- Col. 32, line 50, "KpnI/BgLI<sup>II</sup>" should be --KpnI/Bgl<sup>II</sup>--
- Col. 33, line 51, "EPG" should be --EPO--
- Col. 34, line 54, "requiring" should be --requiring--
- Col. 35, line 49, "Cingote" should be --Congote--
- Col. 35, line 56, "erythropoietin" should be --erythropoietin--
- Col. 35, line 60, "lablled" should be --labelled--
- Col. 36, line 42, "Table V" should be --Figure 5--
- Col. 37, line 20, please delete "8" and insert a bracket "[" immediately before "Asn<sup>2</sup>"
- Col. 37, line 26, please delete "a"
- Col. 37, line 39, "activlty" should be --activity--
- Col. 37, line 44, "Table VI" should be --Figure 6--
- Col. 38, line 61, "really" should be --readily--

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 4,703,008

DATED : October 27, 1987

Page 8 of 8

INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Col. 39, line 20, "membrances" should be --membranes--

Col. 42, line 3, please insert a bracket "]" after "Arg<sup>166</sup>"  
i.e., Arg<sup>166</sup>]hEPO

**Signed and Sealed this  
Fourteenth Day of June, 1988**

*Attest:*

DONALD J. QUIGG

*Attesting Officer*

*Commissioner of Patents and Trademarks*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 4,703,008  
DATED : October 27, 1987  
INVENTOR(S) : FU-KUEN LIN

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 18, line 54, after "83", please insert

--, deposited with the American Type Culture Collection,  
12301 Parklawn Drive, Rockville, Md., under deposit  
accession No. A.T.C.C. 67545 on October 20, 1987--

Column 21, line 25, after "\u039bHE1"  
please insert

--, deposited with the American Type Culture Collection,  
12301 Parklawn Drive, Rockville, Md., under deposit  
accession No. A.T.C.C. 40381 on October 20, 1987--

**Signed and Sealed this**  
**Fourteenth Day of November, 1989**

*Attest:*

JEFFREY M. SAMUELS

*Attesting Officer*

*Acting Commissioner of Patents and Trademarks*