

# **EXHIBIT 3**

## **Part 3 of 3**

5,547,933

13

allowed for the rapid isolation of three positive clones from within a screening of 1,500,000 phage plaques constituting a human genomic library. This was accomplished through use of the above-noted mixture of 128 20-mer probes together with a second set of 128 17-mer probes based on amino acid analysis of a different continuous sequence of human erythropoietin.

The above-noted illustrative procedures constitute the first known instance of the use of multiple mixed oligonucleotide probes in DNA/DNA hybridization processes directed toward isolation of mammalian genomic clones and the first known instance of the use of a mixture of more than 32 oligonucleotide probes in the isolation of cDNA clones.

Numerous aspects and advantages of the invention will be apparent to those skilled in the art upon consideration of the following detailed description which provides illustrations of the practice of the invention in its presently preferred embodiments.

Reference is made to FIGS. 1 through 21, wherein:

FIG. 1 is a graphic representation of a radioimmunoassay analysis of products of the invention;

FIG. 2 shows vector pDSVL-MkE;

FIG. 3 shows vector pSVgHuEPO.

FIG. 4 shows vector pDSVL-gHuEPO.

FIGS. 5A, 5B and 5C (collectively referred to as FIG. 5) show the sequence of monkey EPO cDNA and the encoded EPO.

FIGS. 6A, 6B, 6C, 6D and 6E (collectively referred to as FIG. 6) show the sequence of human genomic EPO DNA and the encoded EPO.

FIG. 7 shows the sequence of the ECEPO gene.

FIG. 8 shows the sequence of the SCEPO gene.

FIG. 9 shows a comparison of the human and monkey EPO polypeptides.

FIG. 10 shows the ECEPO section 1 oligonucleotides.

FIG. 11 shows section 1 of the ECEPO gene.

FIG. 12 shows the ECEPO section 2 oligonucleotides.

FIG. 13 shows section 2 of the ECEPO gene.

FIG. 14 shows the ECEPO section 3 oligonucleotides.

FIG. 15 shows section 3 of the ECEPO gene.

FIG. 16 shows the SCEPO section 1 oligonucleotides.

FIG. 17 shows section 1 of the SCEPO gene.

FIG. 18 shows the SCEPO section 2 oligonucleotides.

FIG. 19 shows section 2 of the SCEPO gene.

FIG. 20 shows the SCEPO section 3 oligonucleotides.

FIG. 21 shows the section 3 of the SCEPO gene.

#### DETAILED DESCRIPTION

According to the present invention, DNA sequences encoding part or all of the polypeptide sequence of human and monkey species erythropoietin (hereafter, at times, "EPO") have been isolated and characterized. Further, the monkey and human origin DNA has been made the subject of eucaryotic and procaryotic expression providing isolatable quantities of polypeptides displaying biological (e.g., immunological) properties of naturally-occurring EPO as well as both in vivo and in vitro biological activities of EPO.

The DNA of monkey species origins was isolated from a cDNA library constructed with mRNA derived from kidney tissue of a monkey in a chemically induced anemic state and whose serum was immunologically determined to include high levels of EPO compared to normal monkey serum. The isolation of the desired cDNA clones containing EPO encoding DNA was accomplished through use of DNA/DNA colony hybridization employing a pool of 128 mixed, radiolabelled, 20-mer oligonucleotide probes and involved the

14

rapid screening of 200,000 colonies. Design of the oligonucleotide probes was based on amino acid sequence information provided by enzymatic fragmentation and sequencing a small sample of human EPO.

The DNA of human species origins was isolated from a human genomic DNA library. The isolation of clones containing EPO-encoding DNA was accomplished through DNA/DNA plaque hybridization employing the above-noted pool of 128 mixed 20-mer oligonucleotide probes and a second pool of 128 radiolabelled 17 mer probes whose sequences were based on amino acids sequence information obtained from a different enzymatic human EPO fragment.

Positive colonies and plaques were verified by means of dideoxy sequencing of clonal DNA using a subset of 16 sequences within the pool of 20-mer probes and selected clones were subjected to nucleotide sequence analysis resulting in deduction of primary structural conformation of the EPO polypeptides encoded thereby. The deduced polypeptide sequences displayed a high degree of homology to each other and to a partial sequence generated by amino acid analysis of human EPO fragments.

A selected positive monkey cDNA clone and a selected positive human genomic clone were each inserted in a "shuttle" DNA vector which was amplified in *E.coli* and employed to transfect mammalian cells in culture. Cultured growth of transfected host cells resulted in culture medium supernatant preparations estimated to contain as much as 3000 mU of EPO per ml of culture fluid.

The following examples are presented by way of illustration of the invention and are specifically directed to procedures carried out prior to identification of EPO encoding monkey cDNA clones and human genomic clones, to procedures resulting in such identification, and to the sequencing, development of expression systems and immunological verification of EPO expression in such systems.

More particularly, Example 1 is directed to amino acid sequencing of human EPO fragments and construction of mixtures of radiolabelled probes based on the results of this sequencing. Example 2 is generally directed to procedures involved in the identification of positive monkey cDNA clones and thus provides information concerning animal treatment and preliminary radioimmunoassay (RIA) analysis of animal sera. Example 3 is directed to the preparation of the cDNA library, colony hybridization screening and verification of positive clones, DNA sequencing of a positive cDNA clone and the generation of monkey EPO polypeptide primary structural conformation (amino acid sequence) information. Example 4 is directed to procedures involved in the identification of positive human genomic clones and thus provides information concerning the source of the genomic library, plaque hybridization procedures and verification of positive clones. Example 5 is directed to DNA sequencing of a positive genomic clone and the generation of human EPO polypeptide amino acid sequence information including a comparison thereof to the monkey EPO sequence information. Example 6 is directed to procedures for construction of a vector incorporating EPO-encoding DNA derived from a positive monkey cDNA clone, the use of the vector for transfection of COS-1 cells and cultured growth of the transfected cells. Example 7 is directed to procedures for construction of a vector incorporating EPO-encoding DNA derived from a positive human genomic clone, the use of the vector for transfection of COS-1 cells and the cultured growth of the transfected cells. Example 8 is directed to immunoassay procedures performed on media supernatants obtained from the cultured growth of trans-

5,547,933

15

fectected cells according to Examples 6 and 7. Example 9 is directed to in vitro and in vivo biological activity of microbially expressed EPO of Examples 6 and 7.

Example 10 is directed to a development of mammalian host expression systems for monkey species EPO cDNA and human species genomic DNA involving Chinese hamster ovary ("CHO") cells and to the immunological and biological activities of products of these expression systems as well as characterization of such products. Example 11 is directed to the preparation of manufactured genes encoding human

16

analysis revealed that within Fragment No. T35 there existed a series of 7 amino acid residues (Val-Asn-Phe-Tyr-Ala-Trp-Lys) which could be uniquely characterized as encoded for by one of 128 possible DNA sequences spanning 20 base pairs. A first set of 128-mer oligonucleotides was therefore synthesized by standard phosphoramidite methods (See, e.g., Beaucage, et al., *Tetrahedron Letters*, 22, pp. 1859-1862 (1981) on a solid support according to the sequence set out in Table II, below.

TABLE II

Residue -	Val	Asn	Phe	Tyr	Ala	Trp	Lys	
3'	CAA	TTG	AAG	ATG	CGA	ACC	TT	- 5'
	T	A	A	A	T			
	G				G			
	C				C			

species EPO and EPO analogs, which genes include a number of preference codons for expression in *E.coli* and yeast host cells, and to expression systems based thereon. Example 12 relates to the immunological and biological activity profiles of expression products of the systems of Example 11.

EXAMPLE 1

A. Human EPO Fragment Amino Acid Sequencing

Human EPO was isolated from urine and subjected to tryptic digestion resulting in the development and isolation of 17 discrete fragments in quantities approximating 100-150 picomoles.

Fragments were arbitrarily assigned numbers and were analyzed for amino acid sequence by microsequence analysis using a gas phase sequencer (Applied Biosystems) to provide the sequence information set out in Table I, below, wherein single letter codes are employed and "X" designates a residue which was not unambiguously determined.

TABLE I

Fragment No.	Sequence Analysis Result
T4a	A-P-P-R
T4b	G-K-L-K
T9	A-L-G-A-Q-K
T13	V-L-E-R
T16	A-V-S-G-L-R
T18	L-F-R
T21	K-L-F-R
T25	Y-L-L-E-A-K
T26a	L-I-C-D-S-R
T26b	L-Y-T-G-E-A-C-R
T27	T-I-T-A-D-T-F-R
T28	E-A-I-S-P-P-D-A-A-M-A-A-P-L-R
T30	E-A-E-X-I-T-T-G-X-A-E-H-X-S-L N-E-X-I-T-V-P
T31	V-Y-S-N-F-L-R
T33	S-L-T-T-L-L-R
T35	V-N-F-Y-A-W-K
T38	G-Q-A-L-L-V-X-S-S-Q-P-W E-P-L-Q-L-H-V-D-K

B. Design and Construction of Oligonucleotide Probe Mixtures

The amino acid sequences set out in Table I were reviewed in the context of the degeneracy of the genetic code for the purpose of ascertaining whether mixed probe procedures could be applied to DNA/DNA hybridization procedures on cDNA and/or genomic DNA libraries. This

Further analysis revealed that within fragment No. T38 there existed a series of 6 amino acid residues (Gln-Pro-Trp-Glu-Pro-Leu) on the basis of which there could be prepared a pool of 128 mixed oligonucleotide 17-mer probes as set out in Table III, below.

TABLE III

Residue -	Gln	Pro	Trp	Glu	Pro	Leu	
3'	GTT	GGA	ACC	CTT	GGA	GA	- 5'
	C	T		C	T	A	
		G			G		
		C			C		

Oligonucleotide probes were labelled at the 5' end with gamma-<sup>32</sup>P-ATP, 7500-8000 Ci/mmol (ICN) using T<sub>4</sub> polynucleotide kinase (NEN).

EXAMPLE 2

A. Monkey Treatment Procedures

Female Cynomolgus monkeys *Macaca fascicularis* (2.5-3 kg, 1.5-2 years old) were treated subcutaneously with a pH 7.0 solution of phenylhydrazine hydrochloride at a dosage level of 12.5 mg/kg on days 1, 3 and 5. The hematocrit was monitored prior to each injection. On day 7, or whenever the hematocrit level fell below 25% of the initial level, serum and kidneys were harvested after administration of 25 mg/kg doses of ketamine hydrochloride. Harvested materials were immediately frozen in liquid nitrogen and stored at -70° C.

B. RIA for EPO

Radioimmunoassay procedures applied for quantitative detection of EPO in samples were conducted according to the following procedures:

An erythropoietin standard or unknown sample was incubated together with antiserum for two hours at 37° C. After the two hour incubation, the sample tubes were cooled on ice, and <sup>125</sup>I labelled erythropoietin was added, the tubes were incubated at 0° C. for at least 15 more hours. Each assay tube contained 500 µl of incubation mixture consisting of 50 µl of diluted immune sera, 10,000 cpm of <sup>125</sup>I-erythropoietin, 5 µl trasylol and 0-250 µl of either EPO standard or unknown sample, with PBS containing 0.1% BSA making up the remaining volume. The antiserum used was the second test bleed of a rabbit immunized with a 1% pure preparation of human urinary erythropoietin. The final antiserum dilution on the assay was adjusted so that the

5,547,933

17

antibody-bound  $^{125}\text{I}$ -EPO did not exceed 10–20% of the input total counts. In general, this corresponded to a final antiserum dilution of from 1:50,000 to 1:100,000.

The antibody-bound  $^{125}\text{I}$ -erythropoietin was precipitated by the addition of 150  $\mu\text{l}$  Staph A. After a 40 min. incubation, the samples were centrifuged and the pellets were washed two times with 0.75 ml 10 mM Tris-HCl pH 8.2 containing 0.15M NaCl, 2 mM EDTA, and 0.05% Triton X-100. The washed pellets were counted in a gamma counter to determine the percent of  $^{125}\text{I}$ -erythropoietin bound. Counts bound by pre-immune sera were subtracted from all final values to correct for nonspecific precipitation. The erythropoietin content of the unknown samples was determined by comparison to the standard curve.

The above procedure was applied to monkey serum obtained in Part A, above, as well as to the untreated monkey serum. Normal serum levels were assayed to contain approximately 36 mU/ml while treated monkey serum contained from 1000 to 1700 mU/ml.

### EXAMPLE 3

#### A. Monkey cDNA Library Construction

Messenger RNA was isolated from normal and anemic monkey kidneys by the guanidinium thiocyanate procedure of Chirgwin, et al., *Biochemistry*, 18, p. 5294 (1979) and poly (A)<sup>+</sup> mRNA was purified by two runs of oligo(dT)-cellulose column chromatography as described at pp. 197–198 in Maniatis, et al., "Molecular Cloning, A Laboratory Manual" (Cold Spring Harbor Laboratory, Cold Spring, Harbor, N.Y., 1982). The cDNA library was constructed according to a modification of the general procedures of Okayama, et al., *Mol. and Cell. Biol.*, 2, pp. 161–170 (1982). The key features of the presently preferred procedures were as follows: (1) pUC8 was used as the sole vector, cut with PstI and then tailed with oligo dT of 60–80 bases in length; (2) HincII digestion was used to remove the oligo dT tail from one end of the vector; (3) first strand synthesis and oligo dG tailing was carried out according to the published procedure; BamHI digestion was employed to remove the oligo dG tail from one end of the Vector; and (5) replacement of the RNA strand by DNA was in the presence of two linkers (GATCTAAAGACCGTCCCCCCCC and ACGGTCTTTA) in a three-fold molar excess over the oligo dG tailed vector.

#### B. Colony Hybridization Procedures For Screening Monkey cDNA Library

Transformed *E. coli* were spread out at a density of 9000 colonies per 10x10 cm plate on nutrient plates containing 50 micrograms/ml Ampicillin. GeneScreen filters (New England Nuclear Catalog No. NEF-972) were pre-wet on a BHI-CAM plate (Bacto brain heart infusion 37 g/L, Casamino acids 2 g/L and agar 15 g/L, containing 500 micrograms/ml Chloramphenicol) and were used to lift the colonies off the plate. The colonies were grown in the same medium for 12 hours or longer to amplify the plasmid copy numbers. The amplified colonies (colony side up) were treated by serially placing the filters over 2 pieces of Whatman 3 MM paper saturated with each of the following solutions:

- (1) 50 mM glucose—25 mM Tris-HCl (pH 8.0)—10 mM EOTA (pH 8.0) for five minutes;
- (2) 0.5M NaOH for ten minutes; and
- (3) 1.0M Tris-HCl (pH 7.5) for three minutes.

The filters were then air dried in a vacuum over at 80° C. for two hours.

18

The filters were then subjected to Proteinase K digestion through treatment with a solution containing 50 micrograms/ml of the protease enzyme in Buffer K [0.1M Tris-HCl (pH 8.0)—0.15M NaCl—10 mM EDTA (pH 8.2)—0.2% SDS]. Specifically, 5 ml of the solution was added to each filter and the digestion was allowed to proceed at 55° C. for 30 minutes, after which the solution was removed.

The filters were then treated with 4 ml of a prehybridization buffer (5xSSPE—0.5% SDS—100 micrograms/ml SS *E. coli* DNA—5xBFP). The prehybridization treatment was carried out at 55° C., generally for 4 hours or longer, after which the prehybridization buffer was removed.

The hybridization process was carried out in the following manner. To each filter was added 3 ml of hybridization buffer (5xSSPE—0.5% SDS—100 micrograms/ml yeast tRNA) containing 0.025 micromoles of each of the 128 probe sequences of Table II (the total mixture being designated the EPV mixture) and the filters were maintained at 48° C. for 20 hours. This temperature was 2° C. less than the lowest of the calculated dissociation temperatures (Td) determined for any of the probes.

Following hybridization, the filters were washed three times for ten minutes on a shaker with 6xSSC—0.1% SDS at room temperature and washed two to three times with 6xSSC—1% SDS at the hybridization temperature (48° C.).

Autoradiography of the filters revealed seven positive clones among the 200,000 colonies screened.

Initial sequence analysis of one of the putative monkey cDNA clones (designated clone 83) was performed for verification purposes by a modification of the procedure of Wallace, et al., *Gene*, 16, pp. 21–26 (1981). Briefly, plasmid DNA from monkey cDNA clone 83 was linearized by digestion with EgoRI and denatured by heating in a boiling water bath. The nucleotide sequence was determined by the dideoxy method of Sanger, et al., *P.N.A.S. (U.S.A.)*, 74, pp. 5463–5467 (1977). A subset of the EPV mixture of probes consisting of 16 sequences was used as a primer for the sequencing reactions.

#### C. Monkey EPO cDNA Sequencing

Nucleotide sequence analysis of clone 83 was carried out by the procedures of Messing, *Methods in Enzymology*, 101, pp. 20–78 (1983). Set out in Table IV is a preliminary restriction map analysis of the approximately 1600 base pair EcoRI/HindIII cloned fragment of clone 83. Approximate locations of restriction endonuclease enzyme recognition sites are provided in terms of number of bases 3' to the EcoRI site at the 5' end of the fragment. Nucleotide sequencing was carried out by sequencing individual restriction fragments with the intent of matching overlapping fragments. For example, an overlap of sequence information provided by analysis of nucleotides in a restriction fragment designated C113 (Sau3A at ~111/SmaI at ~324) and the reverse order sequencing of a fragment designated C73 (AluI at ~424/BstEII at ~203).

TABLE IV

Restriction Enzyme Recognition Site	Approximate Location(s)
EcoRI	1
Sau3A	111
SmaI	180
BstEII	203
SmaI	324
KpnI	371
RsaI	372
AluI	424
PstI	426



5,547,933

19

TABLE IV-continued

Restriction Enzyme Recognition Site	Approximate Location(s)
AluI	430
HpaI	466
AluI	546
PstI	601
PvuII	604
AluI	605
AluI	782
AluI	788
RsaI	792
PstI	807
AluI	841
AluI	927
NcoI	946
Sau3A	1014
AluI	1072
AluI	1115
AluI	1223
PstI	1301
RsaI	1343
AluI	1384
HindIII	1449
AluI	1450
HindIII	1585

Sequencing of approximately 1342 base pairs (within the region spanning the Sau3A site 3' to the EcoRI site and the HindIII site) and analysis of all possible reading frames has allowed for the development of DNA and amino acid sequence information set out in FIG. 5, comprising portions 5A, 5B and 5C. In the Figure, the putative initial amino acid residue of the amino terminal of mature EPO (as verified by correlation to the previously mentioned sequence analysis of twenty amino terminal residues) is designated by the numeral +1. The presence of a methionine-specifying ATG codon (designated -27) "upstream" of the initial amino terminal alanine residue as the first residue designated for the amino acid sequence of the mature protein is indicative of the likelihood that EPO is initially expressed in the cytoplasm in a precursor form including a 27 amino acid "leader" region which is excised prior to entry of mature EPO into circulation. Potential glycosylation sites within the polypeptide are designated by asterisks. The estimated molecular weight of the translated region was determined to be 21,117 daltons and the M.W. of the 165 residues of the polypeptide constituting mature monkey EPO was determined to be 18,236 daltons.

The polypeptide sequence of FIG. 5 may readily be subjected to analysis for the presence of highly hydrophilic regions and/or secondary conformational characteristics indicative of potentially highly immunogenic regions by, e.g., the methods of Hopp, et al., *P.N.A.S. (U.S.A.)*, 78, pp. 3824-3828 (1981) and Kyte et al., *J.Mol.Biol.*, 157, pp. 105-132 (1982) and/or Chou, et al., *Biochem.*, 13, pp. 222-245 (1974) and *Advances in Enzymology*, 47, pp. 45-47 (1978). Computer-assisted analysis according to the Hopp, et al. method is available by means of a program designated PEP Reference Section 6.7 made available by Intelligenetics, Inc., 124 University Avenue, Palo Alto, Calif.

## EXAMPLE 4

## A. Human Genomic Library

A Ch4A phage-borne human fetal liver genomic library prepared according to the procedures of Lawn, et al., *Cell*, supra was obtained and maintained for use in a plaque hybridization assay.

20

## B. Plaque Hybridization Procedures For Screening Human Genomic Library

Phage particles were lysed and the DNAs were fixed on filters (50,000 plaques per filter) according to the procedures of Woo, *Methods In Enzymology*, 68, pp. 389-395 (1979) except for the use of GeneScreen Plus filters (New England Nuclear Catalog No. NEF-972) and NZYAM plates (NACI, 5 g; MgCl<sub>2</sub>-6H<sub>2</sub>O, 2 g; NZ-Amine A, 10 g; yeast extract, 5 g; casamino acids, 2 g; maltose; 2 g; and agar, 15 g per liter). The air-dried filters were baked at 80° C. for 1 hour and then digested with Proteinase K as described in Example 3, Part B. Prehybridization was carried out with a 1M NaCl—1% SDS buffer for 55° C. for 4 hours or more, after which the buffer was removed. Hybridization and post-hybridization washings were carried out as described in Example 3, Part B. Both the mixture of 128 20-mer probes designated EPY and the mixture of 128 17-mer probes of Table III (designated the EPQ mixture) were employed. Hybridization was carried out at 48° C. using the EPV probe mixture. EPQ probe mixture hybridization was carried out at 46° C.—4 degrees below the lowest calculated Td for members of the mixture. Removal of the hybridized probe for rehybridization was accomplished by boiling with 1×SSC—0.1% SDS for two minutes. Autoradiography of the filters revealed three positive clones (reactive with both probe mixtures) among the 1,500,000 phage plaques screened. Verification of the positive clones as being EPO-encoding was obtained through DNA sequencing and electron micrographic visualization of heteroduplex formation with the monkey cDNA of Example 3. This procedure also gave evidence of multiple introns in the genomic DNA sequence.

## EXAMPLE 5

Nucleotide sequence analysis of one of the positive clones (designated λhE1) was carried out and results obtained to date are set out in FIG. 6, comprising portions 6A, 6B, 6C, 6D and 6E.

In FIG. 6, the initial continuous DNA sequence designates a top strand of 620 bases in what is apparently an untranslated sequence immediately preceding a translated portion of the human EPO gene. More specifically, the sequence appears to comprise the 5' end of the gene which leads up to a translated DNA region coding for the first four amino acids (-27 through -24) of a leader sequence ("presequence"). Four base pairs in the sequence prior to that encoding the beginning of the leader have not yet been unambiguously determined and are therefore designated by an "X". There then follows an intron of about 639 base pairs (439 base pairs of which have been sequenced and the remaining 200 base pairs of which are designated "I.S.") and immediately preceding a codon for glutamic acid which has been designated as residue -23 of the translated polypeptide. The exon sequence immediately following is seen to code for amino acid residues through an alanine residue (designated as the +1 residue of the amino acid sequence of mature human EPO) to the codon specifying threonine at position +26, whereupon there follows a second intron consisting of 256 bases as specifically designated. Following this intron is an exon sequence for amino acid residues 27 through 55 and thereafter a third intron comprising 612 base pairs commences. The subsequent exon codes for residues 56 through 115 of human EPO and there then commences a fourth intron of 134 bases as specified. Following the fourth intron is an exon coding for residue Nos. 116 through 166 and a "stop" codon (TGA). Finally, FIG. 6 identifies a sequence of 568 base pairs in what appears to be an untranslated 3' region

5,547,933

21

of the human EPO gene, two base pairs of which ("X") have not yet been unambiguously sequenced.

FIG. 6 thus serves to identify the primary structural conformation (amino acid sequence) of mature human EPO as including 166 specified amino acid residues (estimated M.W.=18,399). Also revealed in the Figure is the DNA sequence coding for a 27 residue leader sequence along with 5' and 3' DNA sequences which may be significant to promoter/operator functions of the human gene operon. Sites for potential glycosylation of the mature human EPO polypeptide are designated in the Figure by asterisks. It is worthy of note that the specific amino acid sequence of FIG. 6 likely constitutes that of a naturally occurring allelic form of human erythropoietin. Support for this position is found in the results of continued efforts at sequencing of urinary isolates of human erythropoietin which provided the finding that a significant number of erythropoietin molecules therein have a methionine at residue 126 as opposed to a serine as shown in the Figure.

FIG. 9, illustrates the extent of polypeptide sequence homology between human and monkey EPO. In the upper continuous line of the Figure, single letter designations are employed to represent the deduced translated polypeptide sequences of human EPO commencing with residue -27 and the lower continuous line shows the deduced polypeptide sequence of monkey EPO commencing at assigned residue number -27. Asterisks are employed to highlight the sequence homologies. It should be noted that the deduced human and monkey EPO sequences reveal an "additional" lysine (K) residue at (human) position 116. Cross-reference to FIG. 6 indicates that this residue is at the margin of a putative mRNA splice junction in the genomic sequence. Presence of the lysine residue in the human polypeptide sequence was further verified by sequencing of a cDNA human sequence clone predated from mRNA isolated from COS-1 cells transformed with the human genomic DNA in Example 7, *infra*.

#### EXAMPLE 6

The expression system selected for initial attempts at microbial synthesis of isolatable quantities of EPO polypeptide material coded for by the monkey cDNA provided by the procedures of Example 3 was one involving mammalian host cells (i.e., COS-1 cells, A.T.C.C. No. CRL-1650). The cells were transfected with a "shuttle" vector capable of autonomous replication in *E. coli* host (by virtue of the presence of pBR322-derived DNA) and the mammalian hosts (by virtue of the presence of SV40 virus-derived DNA).

More specifically, an expression vector was constructed according to the following procedures. The plasmid clone 83 provided in Example 3 was amplified in *E. coli* and the approximately 1.4 kb monkey EPO-encoding DNA was isolated by EcoRI and HindIII digestion. Separately isolated was an approximately 4.0 kb, HindIII/SalI fragment from pBR322. An approximately 30 bp, EcoRI/SalI "linker" fragment was obtained from M13mp10 RF DNA (P and L Laboratories). This linker included, in series, an EcoRI sticky end, followed by SstI, SmaI, BamHI and XbaI recognition sites and a SalI sticky end. The above three fragments were ligated to provide an approximately 5.4 kb intermediate plasmid ("pERS") wherein the EPO DNA was flanked on one side by a "bank" of useful restriction endonuclease recognition sites. pERS was then digested with HindIII and SalI to yield the EPO DNA and the coRI to SalI

22

(M13mp10) linker. The 1.4 kb fragment was ligated with an approximately 4.0 kb BamHI/SalI of pBR322 and another M13mp10 HindIII/BamHI RF fragment linker also having approximately 30 bp. The M13 linker fragment was characterized by a HindIII sticky end, followed by PstI, SalI, XbaI recognition sites and a BamHI sticky end. The ligation product was, again, a useful intermediate plasmid ("pBR-EPO") including the EPO DNA flanked on both sides by banks of restriction site.

The vector chosen for expression of the EPO DNA in COS-1 cells ("pDSVL1") had previously been constructed to allow for selection and autonomous replication in *E. coli*. These characteristics are provided by the origin of replication and Ampicillin resistance gene DNA sequences present in the region spanning nucleotides 2448 through 4362 of pBR322. This sequence was structurally modified by the addition of a linker providing a HindIII recognition immediately adjacent nucleotide 2448 prior to incorporation into the vector. Among the selected vector's other useful properties was the capacity to autonomously replicate in COS-1 cells and the presence of a viral promoter sequence functional in mammalian cells. These characteristics are provided by the origin of replication DNA sequence and "late gene" viral promoter DNA sequence present in the 342 bp sequence spanning nucleotide numbers 5171 through 270 of the SV40 genome. A unique restriction site (BamHI) was provided in the vector and immediately adjacent the viral promoter sequence through use of a commercially available linker sequence (Collaborative Research). Also incorporated in the vector was a 237 base pair sequence (derived as nucleotide numbers 2553 through 2770 of SV40) containing the "late gene" viral mRNA polyadenylation signal (commonly referred to as a transcription terminator). This fragment was positioned in the vector in the proper orientation vis-a-vis the "late gene" viral promoter via the unique BamHI site. Also present in the vector was another mammalian gene at a location not material to potential transcription of a gene inserted at the unique BamHI site, between the viral promoter and terminator sequences. [The mammalian gene comprised an approximately 2,500 bp mouse dihydrofolate reductase (DHFR) minigene isolated from plasmid pMG-1 as in Gasser, et al., *P.N.A.S. (U.S.A.)*, 79, pp. 6522-6526, (1982).] Again, the major operative components of plasmid pDSVL1 comprise nucleotides 2448 through 4362 of pBR322 along with nucleotides 5171 through 270 (342 bp) and 2553 through 2770 (237 bp) of SV40 DNA.

Following procedures described, e.g., in Maniatis, et al., *suu*, the EPO-encoding DNA was isolated from plasmid pBR-EPO as a BamHI fragment and ligated into plasmid pDSVL1 cut with BamHI. Restriction enzyme analysis was employed to confirm insertion of the EPO gene in the correct orientation in two of the resulting cloned vectors (duplicate vectors H and L). See FIG. 2, illustrating plasmid pDSVL-MkE. Vectors with EPO genes in the wrong orientation (vectors F, X, and G) were saved for use as negative controls in transfection experiments designed to determine EPO expression levels in hosts transformed with vectors having EPO DNA in the correct orientation.

Vectors H, L, F, X and G were combined with carrier DNA (mouse liver and spleen DNA) were employed to transfect duplicate 60 mm plates by calcium phosphate microprecipitate methods. Duplicate 60 mm plates were also transfected with carrier DNA as a "mock" transformation negative control. After five days all culture media were tested for the presence of polypeptides possessing the immunological properties of naturally-occurring EPO.

5,547,933

23

## EXAMPLE 7

## A. Initial EPO Expression System Involving COS-1 Cells

The system selected for initial attempts at microbial synthesis of isolatable quantities of human EPO polypeptide material coded for by the human genomic DNA EPO clone, also involved expression in mammalian host cells (i.e., COS-1 cells, A.T.C.C. No. CRL-1650). The human EPO gene was first sub-cloned into a "shuttle" vector which is capable of autonomous replication in both *E. coli* hosts (by virtue of the presence of pBR322 derived DNA) and in the mammalian cell line COS-1 (by virtue of the presence of SV40 virus derived DNA). The shuttle vector, containing the EPO gene, was then transfected into COS-1 cells. EPO polypeptide material was produced in the transfected cells and secreted into the cell culture media.

More specifically, an expression vector was constructed according to the following procedures. DNA isolated from lambda clone  $\lambda$ hE1, containing the human genomic EPO gene, was digested with BamHI and HindIII restriction endonucleases, and a 5.6 Kb DNA fragment known to contain the entire EPO gene was isolated. This fragment was mixed and ligated with the bacterial plasmid pUC8 (Bethesda Research Laboratories, Inc.) which had been similarly digested, creating the intermediate plasmid "pUC8-HuE", providing a convenient source of this restriction fragment.

The vector chosen for expression of the EPO DNA in COS-1 cells (pSV4SEt) had previously been constructed. Plasmid pSV4SEt contained DNA sequences allowing selection and autonomous replication in *E. coli*. These characteristics are provided by the origin of replication and Ampicillin resistance gene DNA sequences present in the region spanning nucleotides 2448 through 4362 of the bacterial plasmid pBR322. This sequence was structurally modified by the addition of a linker providing a HindIII recognition site immediately adjacent to nucleotide 2448. Plasmid pSV4SEt was also capable of autonomous replication in COS-1 cells. This characteristic was provided by a 342 bp fragment containing the SV40 virus origin of replication (nucleotide numbers 5171 through 270). This fragment had been modified by the addition of a linker providing an EcoRI recognition site adjacent to nucleotide 270 and a linker providing a Sall recognition site adjacent nucleotide 5171. A 1061 bp fragment of SV40 was also present in this vector (nucleotide numbers 1711 through 2772 plus a linker providing a Sall recognition site next to nucleotide number 2772). Within this fragment wBs an unique BamHI recognition sequence. In summary, plasmid pSV4SEt contained unique BamHI and HindIII recognition sites, allowing insertion of the human EPO gene, sequences allowing replication and selection in *E. coli*, and sequences allowing replication in COS-1 cells.

In order to insert the EPO gene into pSV4SEt, plasmid pUC8-HuE was digested with BamHI and HindIII restriction endonucleases and the 5.6 kb EPO encoding DNA fragment isolated. pSV4SEt was also digested with BamHI and HindIII and the major 2513 bp fragment isolated (preserving all necessary functions). These fragments were mixed and ligated, creating the final vector "pSVgHuEPO". (See, FIG. 3.) This vector was propagated in *E. coli* and vector DNA isolated. Restriction enzyme analysis was employed to confirm insertion of the EPO gene.

Plasmid pSVgHuEPO DNA was used to express human EPO polypeptide material in COS-1 cells. More specifically, pSVgHuEPO DNA was combined with carrier DNA and transfected into triplicate 60 mm plates of COS-1 cells. As a control, carrier DNA alone was also transfected into

24

COS-1 cells. Cell culture media were sampled five and seven days later and tested for the presence of polypeptides possessing the immunological properties of naturally occurring human EPO.

## B. Second EPO Expression System Involving COS-1 Cells

Still another system was designed to provide improved production of human EPO polypeptide material coded by the human genomic DNA EPO clone in COS-1 cells (A.T.C.C. No. CRL-1650).

In the immediately preceding system, EPO was expressed in COS-1 cells using its own promoter which is within the 5.6 Kb BamHI to HindIII restriction fragment. In the following construction, the EPO gene is altered so that it is expressed using the SV40 late promoter.

More specifically, the cloned 5.6 Kb BamHI to HindIII genomic human EPO restriction fragment was modified by the following procedures. Plasmid pUC8-HuE, as described above, was cleaved with BamHI and with BstEII restriction endonucleases. BstEII cleaves within the 5.6 Kb EPO gene at a position which is 44 base pairs 5' to the initiating ATG coding for the pre-peptide and approximately 680 base pairs 3' to the HindIII restriction site. The approximately 4900 base pair fragment was isolated. A synthetic linker DNA fragment, containing Sall and BstEII sticky ends and an internal BamHI recognition site was synthesized and purified. The two fragments were mixed and ligated with plasmid pBR322 which had been cut with Sall and BamHI to produce the intermediate plasmid pBRgHE. The genomic human EPO gene can be isolated therefrom as a 4900 base pair BamHI digestion fragment carrying the complete structural gene with a single ATG 44 base pairs 3' to BamHI site adjacent the amino terminal coding region.

This fragment was isolated and inserted as a BamHI fragment into BamHI cleaved expression vector plasmid pDSVL1 (described in Example 6). The resulting plasmid, pSVLgHuEPO, as illustrated in FIG. 4, was used to express EPO polypeptide material from COS-1 cells, as described in Examples 6 and 7A.

## EXAMPLE 8

Culture media from growth of the six transfected COS-1 cultures of Example 6 were analyzed by radioimmunoassay according to the procedures set forth in Example 2, Part B. Each sample was assayed at 250, 125, 50, and 25 microliter aliquot levels. Supernatants from growth of cells mock transfected or transfected with vectors having incorrect EPO gene orientation were unambiguously negative for EPO immunoreactivity. For each sample of the two supernatants derived from growth of COS-1 cells transfected with vectors H and L having the EPO DNA in the correct orientation, the % inhibition of  $^{125}$ I-EPO binding to antibody ranged from 72 to 88%, which places all values at the top of the standard curve. The exact concentration of EPO in the culture supernatant could not then reliably be estimated. A quite conservative estimate of 300 mU/ml was made, however, from the value calculation of the largest aliquot size (250 microliter).

A representative culture fluid according to Example 6 and five and seven day culture fluids obtained according to Example 7A were tested in the RIA in order to compare activity of recombinant monkey and human materials to a naturally-occurring human EPO standard and the results are set out in graphic form in FIG. 1. Briefly, the results expectedly revealed that the recombinant monkey EPO significantly competed for anti-human EPO antibody although it was not able to completely inhibit binding under the test conditions. The maximum percent inhibition values



5,547,933

25

for recombinant human EPO however, closely approximated those of the human EPO standard. The parallel nature of the dose response curves suggests immunological identity of the sequences (epitopes) in common. Prior estimates of monkey EPO in culture fluids were re-evaluated at these higher dilution levels and were found to range from 2.91 to 3.12 U/ml. Estimated human EPO production levels were correspondingly set at 392 mU/ml for the five-day growth sample and 567 mU/ml for the seven day growth sample. Estimated monkey EPO production levels in the Example 7B expression system were on the same order or better.

## EXAMPLE 9

Culture fluids prepared according to Examples 6 and 7 were subjected to an in vitro assay for EPO activity according to the procedure of Goldwasser, et al., *Endocrinology*, 97, pp. 315-323 (1975). Estimated monkey EPO values for culture fluids tested ranged from 3.2 to 4.3 U/ml. Human EPO culture fluids were also active in this in vitro assay and, further, this activity could be neutralized by anti-EPO antibody. The recombinant monkey EPO culture fluids according to Example 6 were also subjected to an assay for in vivo biological activity according to the general procedures of Cotes, et al., *Nature*, 191, pp. 1065-1067 (1961) and Hammond, et al., *Ann.N.Y.Acad.Sci.*, 149, pp. 516-527 (1968) and activity levels ranged from 0.94 to 1.24 U/ml.

## EXAMPLE 10

In the previous examples, recombinant monkey or human EPO material was produced from vectors used to transfect COS-1 cells. These vectors replicate in COS-1 cells due to the presence of SV40 T antigen within the cell and an SV40 origin of replication on the vectors. Though these vectors produce useful quantities of EPO in COS-1 cells, expression is only transient (7 to 14 days) due to the eventual loss of the vector. Additionally, only a small percentage of COS-1 became productively transfected with the vectors. The present example describes expression systems employing Chinese hamster ovary (CHO) DHFR<sup>-</sup> cells and the selectable marker, DHFR. [For discussion of related expression systems, see U.S. Pat. No. 4,399,216 and European Patent Applications 117058, 117059 and 117060, all published Aug. 29, 1984.]

CHO DHFR<sup>-</sup> cells (DuX-Bll) CHO K1 cells, Urlaub, et al., *Proc. Nat. Acad. Sci. (U.S.A.)*, Vol. 77, 4461 (1980) lack the enzyme dihydrofolate reductase (DHFR) due to mutations in the structural genes and therefore require the presence of glycine, hypoxanthine, and thymidine in the culture media. Plasmids pDSVL-MkE (Example 6) or pDSVL-gHuEPO (Example 7B) were transfected along with carrier DNA into CHO DHFR<sup>-</sup> cells growing in media containing hypoxanthine, thymidine, and glycine in 60 mm culture plates. Plasmid pSVgHuEPO (Example 7A) was mixed with the plasmid pMG2 containing a mouse dihydrofolate reductase gene cloned into the bacterial plasmid vector pBR322 (per Gasser, et al., supra.) The plasmid mixture and carrier DNA was transfected into CHO DHFR<sup>-</sup> cells. (Cells which acquire one plasmid will generally also acquire a second plasmid). After three days, the cells were dispersed by trypsinization into several 100 mm culture plates in media lacking hypoxanthine and thymidine. Only those cells which have been stably transformed with the DHFR gene, and thereby the EPO gene, survive in this media. After 7-21 days, colonies of surviving cells became apparent. These transformant colonies, after dispersion by trypsinization can

26

be continuously propagated in media lacking hypoxanthine and thymidine, creating new cell strains (e.g., CHO pDSVL-MkEPO, CHO pSVgHuEPO, CHO-pDSVL-gHuEPO).

Culture fluids from the above cell strains were tested in the RIA for the presence of recombinant monkey or human EPO. Media for strain CHO pDSVL-kEPO contained EPO with immunological properties like that obtained from COS-1 cells transfected with plasmid pDSVL-kEPO. A representative 65 hour culture fluid contained monkey EPO at 0.60 U/ml.

Culture fluids from CHO pSVgHuEPO and CHO pDSVL-gHuEPO contained recombinant human EPO with immunological properties like that obtained with COS-1 cells transfected with plasmid pSVgHuEPO or pOSVL-gHuEPO. A representative 3 day culture fluid from CHO pSVgHuEPO contained 2.99 U/ml of human EPO and a 5.5 day sample from CHO pDSVL-gHuEPO had 18.2 U/ml of human EPO as measured by the RIA.

The quantity of EPO produced by the cell strains described above can be increased by gene amplification giving new cell strains of greater productivity. The enzyme dihydrofolate reductase (DHFR) which is the product coded for by the DHFR gene can be inhibited by the drug methotrexate (MTX). More specifically, cells propagated in media lacking hypoxanthine and thymidine are inhibited or killed by MTX. Under the appropriate conditions, (e.g., minimal concentrations of MTX) cells resistant to and able to grow in MTX can be obtained. These cells are found to be resistant to NTX due to an amplification of the number of their OHFR genes, resulting in increased production of OHFR enzyme. The surviving cells can, in turn, be treated with increasing concentrations of NTX, resulting in cell strains containing greater numbers of DHFR genes. "Passenger genes" (e.g., EPO) carried on the expression vector along with the OHFR gene or transformed with the OHFR gene are frequently found also to be increased in their gene copy number.

As examples of practice of this amplification system, cell strain CHO pOSVL-NkE was subjected to increasing NTX concentrations (0 nM, 30 nM and 100 nM). Representative 65-hour culture media samples from each amplification step were assayed by RIA and determined to contain 0.60, 2.45 and 6.10 U/ml, respectively. Cell strain CHO pDSVL-gHuEPO was subjected to a series of increasing NTX concentrations of 30 nM, 50 nM, 100 nM, 200 nM, 1  $\mu$ M, and 5  $\mu$ M MTX. A representative 3-day culture media sample from the 100 nM MTX step contained human EPO at 3089 $\pm$ 129 u/ml as judged by RIA. Representative 48 hour cultural medium samples from the 100 nM and 1  $\mu$ M MTX steps contained, respectively, human EPO at 466 and 1352 U/ml as judged by RIA (average of triplicate assays). In these procedures, 1 $\times$ 10<sup>6</sup> cells were plated in 5 ml of media in 60 mm culture dishes. Twenty-four hours later the media were removed and replaced with 5 ml of serum-free media (high glucose DMEM supplemented with 0.1 mM non-essential amino acids and L-glutamine). EPO was allowed to accumulate for 48 hours in the serum-free media. The media was collected for RIA assay and the cells were trypsinized and counted. The average RIA values of 467 U/ml and 1352 U/ml for cells grown at 100 nM and 1  $\mu$ M MTX, respectively, provided actual yields of 2335 U/plate and 6750 U/plate. The average cell numbers per plate were 1.94 $\times$ 10<sup>6</sup> and 3.12 $\times$ 10<sup>6</sup> cells, respectively. The effective production rates for these culture conditions were thus 1264 and 2167 U/10<sup>6</sup> cells/48 hours.

The cells in the cultures described immediately above are a genetically heterogeneous population. Standard screening



5,547,933

27

procedures are being employed in an attempt to isolate genetically homogeneous clones with the highest production capacity. See, Section A, Part 2, of "Points to Consider in the Characterization of Cell Lines Used to Produce Biologics", Jun. 1, 1984, Office of Biologics Research Review, Center for Drugs and Biologics, U.S. Food and Drug Administration.

The productivity of the EPO producing CHO cell lines described above can be improved by appropriate cell culture techniques. The propagation of mammalian cells in culture generally requires the presence of serum in the growth media. A method for production of erythropoietin from CHO cells in media that does not contain serum greatly facilitates the purification of erythropoietin from the culture medium. The method described below is capable of economically producing erythropoietin in serum-free media in large quantities sufficient for production.

Strain CHO pDSYL-gHuEPO cells, grown in standard cell culture conditions, are used to seed spinner cell culture flasks. The cells are propagated as a suspension cell line in the spinner cell culture flask in media consisting of a 50-50 mixture of high glucose DMEM and Ham's F12 supplemented with 5% fetal calf serum, L-glutamine, Penicillin and Streptomycin, 0.05 mM non-essential amino acids and the appropriate concentration of methotrexate. Suspension cell culture allows the EPO-producing CHO cells to be expanded easily to large volumes. CHO cells, grown in suspension, are used to seed roller bottles at an initial seeding density of  $1.5 \times 10^7$  viable cells per 850 cm<sup>2</sup> roller bottle in 200 ml of media. The cells are allowed to grow to confluency as an adherent cell line over a three-day period. The media used for this phase of the growth is the same as use for growth in suspension. At the end of the three-day growth period, the serum containing media is removed and replaced with 100 ml of serum-free media; 50-50 mixture of high glucose DMEM and Ham's F12 supplemented with 0.05 mM non-essential amino acids and L-glutamine. The roller bottles are returned to the roller bottle incubator for a period of 1-3 hours and the media again is removed and replaced with 100 ml of fresh serum-free media. The 1-3 hour incubation of the serum-free media reduces the concentration of contaminating serum proteins. The roller bottles are returned to the incubator for seven days during which erythropoietin accumulates in the serum-free culture media. At the end of the seven-day production phase, the conditioned media is removed and replaced with fresh serum-free medium for a second production cycle. As an example of the practice of this production system, a representative seven-day, serum-free media sample contained human erythropoietin at  $3892 \pm 409$  U/ml as judged by the RIA. Based on an estimated cell density of 0.9 to  $1.8 \times 10^5$  cells/cm<sup>2</sup>, each 850 cm<sup>2</sup> roller bottle contained from 0.75 to  $1.5 \times 10^8$  cells and thus the rate of production of EPO in the 7-day, 100 ml culture was 750 to 1470 U/10<sup>6</sup> cells/48 hours.

Culture fluids from cell strain CHO pDSVL-MkEPO carried in 10 nM MTX were subjected to RIA in vitro and in vivo EPO activity assays. The conditioned media sample contained  $41.2 \pm 1.4$  U/ml of MkEPO as measured by the RIA,  $41.2 \pm 0.064$  U/ml as measured by the in vitro biological activity assay and  $42.5 \pm 5$  U/ml as measured by the in vivo biological activity assay. Amino acid sequencing of polypeptide products revealed the presence of EPO products, a principle species having 3 residues of the "leader" sequence adjacent the putative amino terminal alanine. Whether this is the result of incorrect membrane processing of the polypeptide in CHO cells or reflects a difference in structure of the amino terminus of monkey ERO vis-a-vis human EPO, is presently unknown.

28

Culture fluids from cell strain CHO pDSVL-gHuEPO were subjected to the three assays. A 5.5 day sample contained recombinant human EPO in the media at a level of 18.2 U/ml by RIA assay,  $15.8 \pm 4.6$  U/ml by in vitro assay and  $16.8 \pm 3.0$  U/ml by in vivo assay.

Culture fluid from CHO pDSVL-GHuEPO cells prepared amplified by stepwise 100 nM TX were subjected to the three assays. A 3.0 day sample contained recombinant human EPO at a level of  $3089 \pm 129$  U/ml by RIA,  $2589 \pm 71.5$  U/ml by in vitro assay, and  $2040 \pm 160$  U/ml by in vivo assay. Amino acid sequencing of this product reveals an amino terminal corresponding to that designated in FIG. 6.

Cell conditioned media from CHO cells transfected with plasmid pDSVL-MkE in 10 nM MTX were pooled, and the MTX dialyzed out over several days, resulting in media with an EPO activity of  $221 \pm 5.1$  U/ml (EPO-CCM). To determine the in vivo effect of the EPO-CCM upon hematocrit levels in normal Balb/C mice, the following experiment was conducted. Cell conditioned media from untransfected CHO cells (CCM) and EPO-CCM were adjusted with PBS. CCM was used for the control group (3 mice) and two dose levels of EPO-CCM—4 units per injection and 44 units per injection—were employed for the experimental groups (2 mice/group). Over the course of 5 weeks, the seven mice were injected intraperitoneally, 3 times per week. After the eighth injection, average hematocrit values for the control group were determined to be 50.4%; for the 4 U group, 55.1%; and, for the 44 U group, 67.9%.

Mammalian cell expression products may be readily recovered in substantially purified form from culture media using HPLC (C<sub>4</sub>) employing an ethanol gradient, preferably at pH7.

A preliminary attempt was made to characterize recombinant glycoprotein products from conditioned medium of COS-1 and CHO cell expression of the human EPO gene in comparison to human urinary EPO isolates using both Western blot analysis and SDS-PAGE. These studies indicated that the CHO-produced EPO material had a somewhat higher molecular weight than the COS-1 expression product which, in turn, was slightly larger than the pooled source human urinary extract. All products were somewhat heterogeneous. Neuraminidase enzyme treatment to remove sialic acid resulted in COS-1 and CHO recombinant products of approximately equal molecular weight which were both nonetheless larger than the resulting asialo human urinary extract. Endoglycosidase F enzyme (EC 3.2.1) treatment of the recombinant CHO product and the urinary extract product (to totally remove carbohydrate from both) resulted in substantially homogeneous products having essentially identical molecular weight characteristics.

Purified human urinary EPO and a recombinant, CHO cell-produced, EPO according to the invention were subjected to carbohydrate analysis according to the procedure of Ledeen, et al. *Methods in Enzymology*, 83(Part D), 139-191 (1982) as modified through use of the hydrolysis procedures of Nesser, et al., *Anal. Biochem.*, 142, 58-67 (1984). Experimentally determined carbohydrate constitution values (expressed as molar ratios of carbohydrate in the product) for the urinary isolate were as follows: Hexoses, 1.73; N-acetylglucosamine, 1; N-acetylneuraminic acid, 0.93; Fucose, 0; and N-acetylgalactosamine, 0. Corresponding values for the recombinant product (derived from CHO pDSVL-gHuEPO 3-day culture media at 100 nM MTX) were as follows: Hexoses, 15.09; N-acetylglucosamine, 1; N-acetylneuraminic acid, 0.998; Fucose, 0; and N-acetylgalactosamine, 0. These findings are consistent with the Western blot and SDS-PAGE analysis described above.

5,547,933

29

Glycoprotein products provided by the present invention are thus comprehensive of products having a primary structural conformation sufficiently duplicative of that of a naturally-occurring erythropoietin to allow possession of one or more of the biological properties thereof and having an average carbohydrate composition which differs from that of naturally-occurring erythropoietin.

## EXAMPLE 11

The present example relates to the total manufacture by assembly of nucleotide bases of two structural genes encoding the human species EPO sequence of FIG. 6 and incorporating, respectively "preferred" codons for expression in *E. coli* and yeast (*S. cerevisiae*) cells. Also described is the construction of genes encoding analogs of human EPO. Briefly stated, the protocol employed was generally as set out in the previously noted disclosure of Alton, et al. (WO 83/04053). The genes were designed for initial assembly of component oligonucleotides into multiple duplexes which, in turn, were assembled into three discrete sections. These sections were designed for ready amplification and, upon removal from the amplification system, could be assembled sequentially or through a multiple fragment ligation in a suitable expression vector.

FIGS. 10 through 15 and 7 illustrate the design and assembly of a manufactured gene encoding a human EPO translation product lacking any leader or presequence but including an initial methionine residue at position -1. Moreover, the gene incorporated in substantial part *E. coli* preference codons and the construction was therefore referred to as the "ECEPO" gene.

More particularly, FIG. 10 illustrates oligonucleotides employed to generate the Section 1 of the ECEPO gene encoding amino terminal residues of the human species polypeptide. Oligonucleotides were assembled into duplexes (1 and 2, 3 and 4, etc.) and the duplexes were then ligated to provide ECEPO Section 1 as in FIG. 11. Note that the assembled section includes respective terminal EcoRI and BamHI sticky ends, that "downstream" of the EcoRI sticky end is a XbaI restriction enzyme recognition site; and that "upstream" of the BamHI sticky end is a KpnI recognition site. Section 1 could readily be amplified using the M13 phage vector employed for verification of sequence of

30

ECEPO Section 2 was constructed with EcoRI and BamHI sticky ends and could be isolated as a KpnI/BglII fragment. Similarly, ECEPO Section 3 was prepared with BamHI and SalI sticky ends and could be isolated from phage RF DNA as a BglII/SalI fragment. The three sections thus prepared can readily be assembled into a continuous DNA sequence (FIG. 7) encoding the entire human species EPO polypeptide with an amino terminal methionine codon (ATG) for *E. coli* translation initiation. Note also that "upstream" of the initial ATG is a series of base pairs substantially duplicating the ribosome binding site sequence of the highly expressed OMP-f gene of *E. coli*.

Any suitable expression vector may be employed to carry the ECEPO. The particular vector chosen for expression of the ECEPO gene as the "temperature sensitive" plasmid pCFM536—a derivative of plasmid pCFM414 (A.T.C.C. 40076)—as described in co-pending U.S. patent application Ser. No. 636,727, filed Aug. 6, 1984, (Published EPO Application No. 136,490) by Charles F. Morris. More specifically, pCFM536 was digested with XbaI and HindIII; the large fragment was isolated and employed in a two-part ligation with the ECEPO gene. Sections 1 (XbaI/KpnI), 2 (KpnI/BglII) and 3 (BglII/SalI) had previously been assembled in the correct order in M13 and the EPO gene was isolated therefrom as a single XbaI/HindIII fragment. This fragment included a portion of the polylinker from M13 mp9 phage spanning the SalI to HindIII sites therein. Control of expression in the resulting expression plasmid, p536, was by means of a lambda P<sub>L</sub> promoter, which itself may be under control of the C<sub>1857</sub> repressor gene (such as provided in *E. coli* strain K12ΔHtrp).

The manufactured ECEPO gene above may be variously modified to encode erythropoietin analogs such as [Asn<sup>2</sup>, des-Pro<sup>2</sup> through Ile<sup>6</sup>]hEPO and [His<sup>7</sup>]hEPO, as described below.

A. [Asn<sup>2</sup>, des-Pro<sup>2</sup> through Ile<sup>6</sup>]hEPO

Plasmid 536 carrying the ECEPO manufactured gene of FIG. 7 as a XbaI to HindIII insert was digested with HindIII and XhoI. The latter endonuclease cuts the ECEPO gene at a unique, 6 base pair recognition site spanning the last base of the codon encoding Asp<sup>8</sup> through the second base of the Arg<sup>10</sup> codon. A XbaI/XhoI "linker" sequence was manufactured having the following sequence:

XbaI	+1	2	7	8	9	XhoI
5'-CTAG	Met	Ala	Asn	Cys	Asp	
3'-	ATG	GCT	AAT	TGC	GAC-	3'
	TAC	CGA	TTA	ACG	CTG	AGCT-5'

the section. Some difficulties were encountered in isolating the section as an XbaI/KpnI fragment from RF DNA generated in *E. coli*, likely due to methylation of the KEqI recognition site bases within the host. Single-stranded phage DNA was therefore isolated and rendered into double-stranded form in vitro by primer extension and the desired double-stranded fragment was thereafter readily isolated.

ECEPO gene Sections 2 and 3 FIGS. 13 and 15 were constructed in a similar manner from the oligonucleotides of FIGS. 12 and 14, respectively. Each section was amplified in the M13 vector employed for sequence verification and was isolated from phage DNA. As is apparent from FIG. 13,

The XbaI/XhoI linker and the XhoI/HindIII ECEPO gene sequence fragment were inserted into the large fragment resulting from XbaI and HindIII digestion of plasmid pCFM526—a derivative of plasmid pCFM414 (A.T.C.C. 40076)—as described in co-pending U.S. patent application Ser. No. 636,727, filed Aug. 6, 1984, by Charles F. Morris, to generate a plasmid-borne DNA sequence encoding *E. coli* expression of the Met<sup>-1</sup> form of the desired analog.

B. [His<sup>7</sup>]hEPO

Plasmid 536 was digested with HindIII and XhoI as in part A above. A XbaI/XhoI linker was manufactured having the following sequence:

5,547,933

31

32

XbaI	+1	2	3	4	5	6	7	8	9	XhoI
5'-CTAG	Met ATG	Ala GCT	Pro CCG	Pro CCA	Arg CGT	Leu CTG	Ile ATC	His CAT	Asp GAC	3'
3'-	TAC	CGA	GGC	GGT	GCA	GAC	TAG	GTA	CTG	AGCT-5'

The linker and the XhoI/HindIII ECEPO sequence fragment were then inserted into pCFM526 to generate a plasmid-borne DNA sequence encoding *E. coli* expression of the Met<sup>-1</sup> form of the desired analog.

Construction of a manufactured gene ("SCEPO") incorporating yeast preference codons is as described in the following FIGS. 16 through 21 and 8. As was the case with the ECEPO gene, the entire construction involved formation of three sets of oligonucleotides (FIGS. 16, 18 and 20 which were formed into duplexes and assembled into sections (FIGS. 17, 19 and 21). Note that synthesis was facilitated in part by use of some sub-optimal codons in both the SCEPO and ECEPO constructions, i.e., oligonucleotides 7-12 of Section 1 of both genes were identical, as were oligonucleotides 1-6 of Section 2 in each gene.

The assembled SCEPO sections were sequenced in M13 and Sections 1, 2 and 3 were isolatable from the phage as indIII/KpnI, KpnI/BglII, and BglII/SalI fragments.

The presently preferred expression system for SCEPO gene products is a secretion system based on *S. cerevisiae*  $\alpha$ -factor secretion, as described in copending U.S. patent application Ser. No. 487,753, filed Apr. 22, 1983, by Grant A. Bitter, published Oct. 31, 1984 as European Patent Application 0 123,294. Briefly put, the system involves constructions wherein DNA encoding the leader sequence of the yeast  $\alpha$ -factor gene product is positioned immediately 5' to the coding region of the exogenous gene to be expressed. As a result, the gene product translated includes a leader or signal sequence which is "processed off" by an endogenous yeast enzyme in the course of secretion of the remainder of the product. Because the construction makes use of the  $\alpha$ -factor translation initiation (ATG) codon, there was no need to provide such a codon at the -1 position of the SCEPO gene. As may be noted from FIG. 8, the alanine (+1) encoding sequence is preceded by a linker sequence allowing for direct insertion into a plasmid including the DNA for the first 80 residues of the  $\alpha$ -factor leader following the  $\alpha$ -factor promoter. The specific preferred construction for SCEPO gene expression involved a four-part ligation including the above-noted SCEPO section fragments and the large fragment of HindIII/SalI digestion of plasmid p $\alpha$ C3. From the resulting plasmid p $\alpha$ C3/SCEPO, the  $\alpha$ -factor promoter and leader sequence and SCEPO gene were isolated by digestion with BamHI and ligated into BamHI digested plasmid DYE to form expression plasmid pYE/SCEPO.

#### EXAMPLE 12

The present example relates to expression of recombinant products of the manufactured ECEPO and SCEPO genes within the expression systems of Example 11.

In use of the expression system designed for use of *E. coli* host cells, plasmid p536 of Example 11 was transformed into AM7 *E. coli* cells previously transformed with a suitable plasmid, pMW1, harboring a C<sub>1857</sub> gene. Cultures of cells in LB broth (Ampicillin 50  $\mu$ g/ml and kanamycin 5  $\mu$ g/ml, preferably with 10 mM MgSO<sub>4</sub>) were maintained at 28° C. and upon growth of cells in culture to O.D.<sub>600</sub>=0.1, EPO

expression was induced by raising the culture temperature to 42° C. Cells grown to about 40 O.D. provided EPO production (as estimated by gel) of about 5 mg/OD liter.

Cells were harvested, lysed, broken with French Press (10,000 psi) and treated with lysozyme and NP-40 detergent. The pellet resulting from 24,000xg centrifugation was solubilized with guanidine HCl and subjected to further purification in a single step by means of C<sub>4</sub> (Vydac) Reverse Phase HPLC (EtOH, 0-80%, 50 mM NH<sub>4</sub>Ac, pH 4.5). Protein sequencing revealed the product to be greater than 95% pure and the products obtained revealed two different amino terminals, A-P-P-R . . . and P-P-R . . . in a relative quantitative ratio of about 3 to 1. This latter observation of hEPO and [des Ala<sup>1</sup>]hEPO products indicates that amino terminal "processing" within the host cells serves to remove the terminal methionine and in some instances the initial alanine. Radioimmunoassay activity for the isolates was at a level of 150,000 to 160,000 U/mg; in vitro assay activity was at a level of 30,000 to 62,000 U/mg; and in vivo assay activity ranged from about 120 to 720 U/mg. (Cf., human urinary isolate standard of 70,000 U/mg in each assay.) The dose response curve for the recombinant product in the in vivo assay differed markedly from that of the human urinary EPO standard.

The EPO analog plasmids formed in parts A and B of Example 11 were each transformed into pMW1-transformed AM7 *E. coli* cells and the cells were cultured as above. Purified isolates were tested in both RIA and in vitro assays RIA and in vitro assay values for [Asn<sup>2</sup>, des-Pro<sup>2</sup> through Ile<sup>6</sup>]hEPO expression products were approximately 11,000 U/mg and 6,000 U/mg protein, respectively, while the assay values for [His<sup>7</sup>]hEPO were about 41,000 U/mg and 14,000 U/mg protein, respectively, indicating that the analog products were from one-fourth to one-tenth as "active" as the "parent" expression product in the assays.

In the expression system designed for use of *S. cerevisiae* host cells, plasmid pYE/SCEPO was transformed into two different strains, YSDP4 (genotype  $\alpha$  pep4-3 trpl) and RK81 (genotype pep4-3 trpl). Transformed YSDP4 hosts were grown in SD medium (Methods in Yeast Genetics, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., p. 62 (1983) supplemented with casamino acids at 0.5%, pH 6.5 at 30° C. Media harvested when the cells had been grown to 36 O.D. contained EPO products at levels of about 244 U/ml (97  $\mu$ g/OD liter by RIA). Transformed RK81 cells grown to either 6.5 O.D. or 60 O.D. provided media with EPO concentrations of about 80-90 U/ml C34  $\mu$ g/OD liter by RIA). Preliminary analyses reveal significant heterogeneity in products produced by the expression system, likely to be due to variations in glycosylation of proteins expressed, and relatively high mannose content of the associated carbohydrate.

Plasmids p $\alpha$ C3 and pYE in HB101 *E. coli* cells were deposited in accordance with the Rules of Practice of the U.S. Patent Office on Sep. 27, 1984, with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Md., under deposit numbers A.T.C.C. 39881 and A.T.C.C. 39882, respectively. Plasmids pCFM526 in AM7 cells, pCFM536 in JM103 cells, and pMW1 in JM103 cells were likewise



5,547,933

33

deposited on Nov. 21, 1984 as A.T.C.C. 39932, 39934, and 39933, respectively. *Saccharomyces cerevisiae* strains YSPD4 and RK81 were deposited on Nov. 21, 1984 as A.T.C.C. 20734 and 20733, respectively.

It should be readily apparent from consideration of the above illustrative examples that numerous exceptionally valuable products and processes are provided by the present invention in its many aspects.

Polypeptides provided by the invention are conspicuously useful materials, whether they are microbially expressed products or synthetic products, the primary, secondary or tertiary structural conformation of which was first made known by the present invention.

As previously indicated, recombinant-produced and synthetic products of the invention share, to varying degrees, the in vitro biological activity of EPO isolates from natural sources and consequently are projected to have utility as substitutes for EPO isolates in culture media employed for growth of erythropoietic cells in culture. Similarly, to the extent that polypeptide products of the invention share the in vivo activity of natural EPO isolates they are conspicuously suitable for use in erythropoietin therapy procedures practiced on mammals, including humans, to develop any or all of the effects heretofore attributed in vivo to EPO, e.g., stimulation of reticulocyte response, development of ferrokinetic effects (such as plasma iron turnover effects and marrow transit time effects), erythrocyte mass changes, stimulation of hemoglobin C synthesis (see, Eschbach, et al., supra) and, as indicated in Example 10, increasing hematocrit levels in mammals. Included within the class of humans treatable with products of the invention are patients generally requiring blood transfusions and including trauma victims, surgical patients, renal disease patients including dialysis patients, and patients with a variety of blood composition affecting disorders, such as hemophilia, sickle cell disease, physiologic anemias, and the like. The minimization of the need for transfusion therapy through use of EPO therapy can be expected to result in reduced transmission of infectious agents. Products of the invention, by virtue of their production by recombinant methods, are expected to be free of pyrogens, natural inhibitory substances, and the like, and are thus likely to provide enhanced overall effectiveness in therapeutic processes vis-a-vis naturally derived products. Erythropoietin therapy with products of the present invention is also expected to be useful in the enhancement of oxygen carrying capacity of individuals encountering hypoxic environmental conditions and possibly in providing beneficial cardiovascular effects.

A preferred method for administration of polypeptide products of the invention is by parenteral (e.g., IV, IM, SC, or IP) routes and the compositions administered would ordinarily include therapeutically effective amounts of product in combination with acceptable diluents, carriers and/or adjuvants. Preliminary pharmacokinetic studies indicate a longer half-life in vivo for monkey EPO products when administered IM rather than IV. Effective dosages are expected to vary substantially depending upon the condition treated but therapeutic doses are presently expected to be in the range of 0.1 (-7 U) to 100 (-7000 U) g/kg body weight of the active material. Standard diluents such as human serum albumin are contemplated for pharmaceutical compositions of the invention, as are standard carriers such as saline.

Adjuvant materials suitable for use in compositions of the invention include compounds independently noted for erythropoietic stimulatory effects, such as testosterone, proge-

34

tor cell stimulators, insulin-like growth factor, prostaglandins, serotonin, cyclic AMP, prolactin and triiodothyronine, as well as agents generally employed in treatment of aplastic anemia, such as methenolene, stanozolol and nandrolone [see, e.g., Resegotti, et al., *Panminerva Medica*, 23,, 243-248 (1981); McGonigle, et al., *Kidney Int.*, 25(2), 437-444 (1984); Pavlovic-Kantera, et al., *Expt.Hematol.*, 8(Supp. 8), 283-291 (1980); and Kurtz, *FEBS Letters*, 14a(1), 105-108 (1982)]. Also contemplated as adjuvants are substances reported to enhance the effects of, or synergize, erythropoietin or asialo-EPO, such as the adrenergic agonists, thyroid hormones, androgens and BPA [see, Dunn, "Current Concepts in Erythropoiesis", John Wiley and Sons (Chichester, England, 1983); Welland, et al., *Blut*, 44(3), 173-175 (1982); Kalmanti, *Kidney Int.*, 22, 383-391 (1982) Shahidi, *New.Eng.J.Med.*, 289, 72-80 (1973); Fisher, et al., *Steroids*, 30(6), 833-845 (1977); Urabe, et al., *J.Exp.Med.*, 149, 1314-1325 (1979); and Billat, et al., *Expt.Hematol.*, 10(1), 133-140 (1982)] as well as the classes of compounds designated "hepatic erythropoietic factors" [see, Naughton, et al., *Acta.Haemat.*, 69, 171-179 (1983)] and "erythrotropins" [as described by Congote, et al. in Abstract 364, Proceedings 7th International Congress of Endocrinology (Quebec City, Quebec, Jul. 1-7, 1984); Congote, *Biochem. Biophys.Res.Comm.*, 115(2), 447-483 (1983) and Congote, *Anal.Biochem.*, 140, 428-433 (1984)] and "erythrogenins" [(as described in Rothman, et al., *J.Surg.Oncol.*, 20, 105-108 (1982)]. Preliminary screenings designed to measure erythropoietic responses of ex-hypoxic polycythemic mice pre-treated with either 5- $\alpha$ -dihydrotestosterone or nandrolone and then given erythropoietin of the present invention have generated equivocal results.

Diagnostic uses of polypeptides of the invention are similarly extensive and include use in labelled and unlabelled forms in a variety of immunoassay techniques including RIA's ELISA's and the like, as well as a variety of in vitro and in vivo activity assays. See, e.g., Dunn, et al., *Expt.Hematol.*, 11(7), 590-600 (1983); Gibson, et al., *Pathology*, 16, 155-156 (1984); Krystal, *Expt.Hematol.*, 11(7), 649-660 (1983); Saito, et al., *Jap.J.Med.*, 23(1), 16-21 (1984); Nathan, et al., *New Eng.J.Med.*, 308(9), 520-522 (1983); and various references pertaining to assays referred to therein. Polypeptides of the invention, including synthetic peptides comprising sequences of residues of EPO first revealed herein, also provide highly useful pure materials for generating polyclonal antibodies and "banks" of monoclonal antibodies specific for differing continuous and discontinuous epitopes of EPO. As one example, preliminary analysis of the amino acid sequences of FIG. 6 in the context of hydrophobicity according to Hopp, et al., *P.N.A.S. (U.S.A.)*, 78, pp. 3824-3828 (1981) and of secondary structures according to Chou, et al., *Ann.Rev. Biochem.*, 47, p. 251 (1978) revealed that synthetic peptides duplicative of continuous sequences of residues spanning positions 41-57 inclusive, 116-128 inclusive and 144-166 inclusive are likely to produce a highly antigenic response and generate useful monoclonal and polyclonal antibodies immunoreactive with both the synthetic peptide and the entire protein. Such antibodies are expected to be useful in the detection and affinity purification of EPO and EPO-related products.

Illustratively, the following three synthetic peptides were prepared:

- (1) hEPO 41-57, V-P-D-T-K-V-N-F-Y-A-W-K-R-M-E-V-G;
- (2) hEPO 116-128, K-E-A-I-S-P-P-D-A-A-S-A-A;
- (3) hEPO 144-166, V-Y-S-N-F-L-R-G-K-L-K-L-Y-T-G-E-A-C-R-T-G-D-R. Preliminary immunization studies



5,547,933

35

employing the above-noted polypeptides have revealed a relatively weak positive response to hEPO 41-57, no appreciable response to hEPO 116-128, and a strong positive response to hEPO 144-166, as measured by capacity of rabbit serum antibodies to immunoprecipitate <sup>125</sup>I-labelled human urinary EPO isolates. Preliminary in vivo activity studies on the three peptides revealed no significant activity either alone or in combination.

While the deduced sequences of amino acid residues of mammalian EPO provided by the illustrative examples essentially define the primary structural conformation of mature EPO, it will be understood that the specific sequence of 165 amino acid residues of monkey species EPO in FIG. 5 and the 166 residues of human species EPO in FIG. 6 do not limit the scope of useful polypeptides provided by the invention. Comprehended by the present invention are those various naturally-occurring allelic forms of EPO which past research into biologically active mammalian polypeptides such as human  $\gamma$  interferon indicates are likely to exist. (Compare, e.g., the human immune interferon species reported to have an arginine residue at position No. 140 in EPO published application 0 077 670 and the species reported to have glutamine at position No. 140 in Gray, et al., *Nature*, 295, pp. 503-508 (1982). Both species are characterized as constituting "mature" human  $\gamma$  interferon sequences.) Allelic forms of mature EPO polypeptides may vary from each other and from the sequences of FIGS. 5 and 6 in terms of length of sequence and/or in terms of deletions, substitutions, insertions or additions of amino acids in the sequence, with consequent potential variations in the capacity for glycosylation. As noted previously, one putative allelic form of human species EPO is believed to include a methionine residue at position 126. Expectedly, naturally-occurring allelic forms of EPO-encoding DNA genomic and cDNA sequences are also likely to occur which code for the above-noted types of allelic polypeptides or simply employ differing codons for designation of the same polypeptides as specified.

In addition to naturally-occurring allelic forms of mature EPO, the present invention also embraces other "EPO products" such as polypeptide analogs of EPO and fragments of "mature" EPO. Following the procedures of the above-noted published application by Alton, et al. (WO/83/04053) one may readily design and manufacture genes coding for microbial expression of polypeptides having primary conformations which differ from that herein specified for mature EPO in terms of the identity or location of one or more residues (e.g., substitutions, terminal and intermediate additions and deletions). Alternately, modifications of cDNA and genomic EPO genes may be readily accomplished by well-known site-directed mutagenesis techniques and employed to generate analogs and derivatives of EPO. Such EPO products would share at least one of the biological properties of EPO but may differ in others. As examples, projected EPO products of the invention include those which are foreshortened by e.g., deletions [Asn<sup>2</sup>, des-Pro<sup>2</sup> through Ile<sup>6</sup>]hEPO, [des-Thr<sup>163</sup> through Arg<sup>166</sup>]hEPO and "Δ27-55hEPO" the latter having the residues coded for by an entire exon deleted; or which are more stable to hydrolysis (and, therefore, may have more pronounced or longer lasting effects than naturally-occurring EPO); or which have been altered to delete one or more potential sites for glycosylation (which may result in higher activities for yeast-produce products); or which have one or more cystein residues deleted or replaced by, e.g., histidine or serine residues (such as the analog [His<sup>7</sup>]hEPO) and are potentially more easily isolated

36

in active form from microbial systems; or which have one or more tyrosine residues replaced by phenylalanine (such as the analogs [Phe<sup>15</sup>]hEPO, [Phe<sup>49</sup>]hEPO, and [Phe<sup>145</sup>]hEPO) and may bind more or less readily to EPO receptors on target cells. Also comprehended are polypeptide fragments duplicating only a part of the continuous amino acid sequence or secondary conformations within mature EPO, which fragments may possess one activity of EPO (e.g., receptor binding) and not others (e.g., erythropoietic activity). Especially significant in this regard are those potential fragments of EPO which are elucidated upon consideration of the human genomic DNA sequence of FIG. 6, i.e., "fragments" of the total continuous EPO sequence which are delineated by intron sequences and which may constitute distinct "domains" of biological activity. It is noteworthy that the absence of in vivo activity for any one or more of the "EPO products" of the invention is not wholly preclusive of therapeutic utility (see, Welland, et al., supra) or of utility in other contexts, such as may be quite useful in treatment of polycythemias or cases of overproduction of EPO [see, e.g., Adamson, *Hosp.Practice*, 18(12), 49-57 (1983), and Hellmann, et al., *Clin.Lab.Haemat.*, 5, 335-342 (1983)].

According to another aspect of the present invention, the cloned DNA sequences described herein which encode human and monkey EPO polypeptides are conspicuously valuable for the information which they provide concerning the amino acid sequence of mammalian erythropoietin which has heretofore been unavailable despite decades of analytical processing of isolates of naturally-occurring products. The DNA sequences are also conspicuously valuable as products useful in effecting the large scale microbial synthesis of erythropoietin by a variety of recombinant techniques. Put another way, DNA sequences provided by the invention are useful in generating new and useful viral and circular plasmid DNA vectors, new and useful transformed and transfected microbial procaryotic and eucaryotic host cells (including bacterial and yeast cells and mammalian cells grown in culture), and new and useful methods for cultured growth of such microbial host cells capable of expression of EPO and EPO products. DNA sequences of the invention are also conspicuously suitable materials for use as labelled probes in isolating EPO and related protein encoding cDNA and genomic DNA sequences of mammalian species other than human and monkey species herein specifically illustrated. The extent to which DNA sequences of the invention will have use in various alternative methods of protein synthesis (e.g., in insect cells) or in genetic therapy in humans and other mammals cannot yet be calculated. DNA sequences of the invention are expected to be useful in developing transgenic mammalian species which may serve as eucaryotic "hosts" for production of erythropoietin and erythropoietin products in quantity. See, generally, Palmiter, et al., *Science*, 222(4625), 809-814 (1983).

Viewed in this light, therefore, the specific disclosures of the illustrative examples are clearly not intended to be limiting upon the scope of the present invention and numerous modifications and variations are expected to occur to those skilled in the art. As one example, while DNA sequences provided by the illustrative examples include cDNA and genomic DNA sequences, because this application provides amino acid sequence information essential to manufacture of DNA sequence, the invention also comprehends such manufactured DNA sequences as may be constructed based on knowledge of EPO amino acid sequences. These may code for EPO (as in Example 12) as well as for EPO fragments and EPO polypeptide analogs (i.e., "EPO Products") which may share one or more biological prop-

5,547,933

37

erties of naturally-occurring EPO but not share others (or possess others to different degrees).

DNA sequences provided by the present invention are thus seen to comprehend all DNA sequences suitable for use in securing expression in a procaryotic or eucaryotic host cell of a polypeptide product having at least a part of the primary structural conformation and one or more of the biological properties of erythropoietin, and selected from among: (a) the DNA sequences set out in FIGS. 5 and 6; (b) DA sequences which hybridize to the DA sequences defined in (a) or fragments thereof; and (c) DNA sequences which, but for the degeneracy of the genetic code, would hybridize to the DNA sequences defined in (a) and (b). It is noteworthy in this regard, for example, that existing allelic monkey and human EPO gene sequences and other mammalian species gene sequences are expected to hybridize to the sequences of FIGS. 5 and 6 or to fragments thereof. Further, but for the degeneracy of the genetic code, the SCEPO and ECEPO genes and the manufactured or mutagenized cDNA or genomic DNA sequences encoding various EPO fragments and analogs would also hybridize to the above-mentioned DNA sequences. Such hybridizations could readily be carried out under the hybridization conditions described herein with respect to the initial isolation of the monkey and human EPO-encoding DNA or more stringent conditions, if desired to reduce background hybridization.

In a like manner, while the above examples illustrate the invention of microbial expression of EPO products in the context of mammalian cell expression of DNA inserted in a hybrid vector of bacterial plasmid and viral genomic origins, a wide variety of expression systems are within the contemplation of the invention. Conspicuously comprehended are expression systems involving vectors of homogeneous origins applied to a variety of bacterial, yeast and mammalian cells in culture as well as to expression systems not involving vectors (such as calcium phosphate transfection of cells). In this regard, it will be understood that expression of, e.g., monkey origin DNA in monkey host cells in culture and human host cells in culture, actually constitute instances of "exogenous" DNA expression inasmuch as the EPO DNA whose high level expression is sought would not have its origins in the genome of the host. Expression systems of the invention further contemplate these practices resulting in cytoplasmic formation of EPO products and accumulation of glycosylated and non-glycosylated EPO products in host cell cytoplasm or membranes (e.g., accumulation in bacterial periplasmic spaces) or in culture medium supernatants as above illustrated, or in rather uncommon systems such as *Paeriginosa* expression systems (described in Gray, et al., *Biotechnology*, 2, pp. 161-165 (1984)).

Improved hybridization methodologies of the invention, while illustratively applied above to DNA/DNA hybridization screenings are equally applicable to RNA/RNA and RNA/DNA screening. Mixed probe techniques as herein illustrated generally constitute a number of improvements in hybridization processes allowing for more rapid and reliable polynucleotide isolations. These many individual processing improvements include: improved colony transfer and maintenance procedures; use of nylon-based filters such as Gene-Screen and GeneScreen Plus to allow reprobing with same filters and repeated use of the filter, application of novel protease treatments [compared, e.g., to Taub, et al. *Anal. Biochem.*, 126, pp. 222-230 (1982)]; use of very low individual concentrations (on the order of 0.025 picomole) of a large number of mixed probes (e.g., numbers in excess of 32); and, performing hybridization and post-hybridization

38

steps under stringent temperatures closely approaching (i.e., within 4° C. and preferably within 2° C. away from) the lowest calculated dissociation temperature of any the mixed probes employed. These improvements combine to provide results which could not be expected to attend their use. This is amply illustrated by the fact that mixed probe procedures involving 4 times the number of probes ever before reported to have been successfully used in even cDNA screens on messenger RNA species of relatively low abundance were successfully applied to the isolation of a unique sequence gene in a genomic library screening of 1,500,000 phage plaques. This feat was accomplished essentially concurrently with the publication of the considered opinion of Anderson, et al., *supra*, that mixed probe screening methods were "... impractical for isolation of mammalian protein genes when corresponding RNA's are unavailable.

What is claimed is:

1. A non-naturally occurring erythropoietin glycoprotein product having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and having glycosylation which differs from that of human urinary erythropoietin.

2. The non-naturally occurring EPO glycoprotein product according to claim 1 wherein said product has a higher molecular weight than human urinary EPO as measured by SDS-PAGE.

3. A non-naturally occurring glycoprotein product of the expression in a mammalian host cell of an exogenous DNA sequence comprising a DNA sequence encoding human erythropoietin said product possessing the *in vivo* biological property of causing bone marrow cells to increase production of reticulocytes and red blood cells.

4. A non-naturally occurring human erythropoietin glycoprotein possessing the *in vivo* biological property of causing bone marrow cells to increase production of reticulocytes and red blood cells which is the product of the process comprising the steps of:

(a) growing, under suitable nutrient conditions, mammalian host cells transformed or transfected with an isolated DNA sequence encoding the human erythropoietin amino acid sequence set out in FIG. 6 or a fragment thereof; and

(b) isolating a glycosylated erythropoietin polypeptide therefrom.

5. A non-naturally occurring human erythropoietin glycoprotein possessing the *in vivo* biological property of causing bone marrow cells to increase production of reticulocytes and red blood cells which is the product of the process comprising the steps of:

(a) growing, under suitable nutrient conditions, mammalian host cells transformed or transfected with an isolated DNA sequence comprising a sequence encoding the leader sequence of human erythropoietin set out in FIG. 6; and

(b) isolating a glycosylated erythropoietin polypeptide therefrom.

6. A non-naturally occurring glycoprotein product of the expression in a non-human eucaryotic host of an exogenous DNA sequence comprising a DNA sequence encoding human erythropoietin, said product possessing the *in vivo* biological property of causing human bone marrow cells to increase production of reticulocytes and red blood cells and having an average carbohydrate composition which differs from that of naturally occurring erythropoietin.

7. The glycoprotein product according to claim 3, 4, 5 or 6 wherein the host cell is a non-human mammalian cell.

8. The glycoprotein product according to claim 7 wherein the non-human mammalian cell is a CHO cell.

5,547,933

39

9. A pharmaceutical composition comprising an effective amount of a glycoprotein product effective for erythropoietin therapy according to claim 1, 2, 3, 4, 5 or 6 and a pharmaceutically acceptable diluent, adjuvant or carrier.

10. A method for providing erythropoietin therapy to a mammal comprising administering an effective amount of a pharmaceutical composition of claim 9.

11. A method for treating a kidney dialysis patient which comprises administering a pharmaceutical composition of claim 9 in an amount effective to increase the hematocrit level of said patient.

12. A pharmaceutical composition comprising an effective amount of a glycoprotein product effective for erythropoi-

40

etin therapy according to claim 7 and a pharmaceutically acceptable diluent, adjuvant or carrier.

13. A method for providing erythropoietin therapy to a mammal comprising administering an effective amount of a pharmaceutical composition of claim 12.

14. A method for treating a kidney dialysis patient which comprises administering a pharmaceutical composition of claim 12 in an amount effective to increase the hematocrit level of said product.

\* \* \* \* \*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

Page 1 of 7

PATENT NO. : 5,547,933  
DATED : August 20, 1996  
INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In the References, page 1, column 2, line 3  
"012394" should be -- 0123294 --.

In Other Publications, page 2, column 1, line 60,  
Chia Reference, "(1992)" should -- (1982) --.

In Other Publications, page 5, column 2, line 38,  
Konwalinka Reference, "Humlan" should be -- Human --.

In Other Publications, page 6, column 1, line 51,  
Lee-Huang Reference, "620, 622 and 624" should be -- 620-624 --.

In Other Publications, page 7, column 1, lines 7-8,  
Mirand Reference, "94, 96, 98, 100, 102, 104, 106" should be -- 94-106 --.

In Other Publications, page 7, column 1, line 16,  
Montgomery Reference, "67-680" should be -- 673-680 --.

In Other Publications, page 9, column 1, line 9,  
Storring Reference, "459-489" should be -- 459-484 --.

In Other Publications, page 9, column 2, line 59, Wallace Reference insert  
(1981) after 3647-3657 but before [D].





UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

Page 3 of 7

PATENT NO. : 5,547,933

DATED : August 20, 1996

INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

column 1, line 50, "uracil" should be -- uracil (U) --.

column, 1, line 62, "sequences" should be -- sequences. --.

column 1, line 62, "'message'" should be -- "message,".

column 4, line 15, "DnAs" should be -- DNAs --.

column 4, line 25, "Wallase" should be -- Wallace --.

column 4, line 25, "(1981;" should be -- (1981); --.

column 4, line 31, "wallace," should be -- Wallace, --.

column 4, line 61, "off" should be -- of --.

column 5, line 18, "amine" should be -- amino --.

column 5, line 31, "amine" should be -- amino --.

column 5, line 36, "amine" should be -- amino --.

column 5, line 50, " $\beta$ and" should be --  $\beta$  and --.

column 8, line 27, "anti bodies" should be -- antibodies --.

column 10, line 21, "arian" should be -- avian --.

column 10, line 46, "100 U" should be -- 100U --.

column 10, line 47, "500 U" should be -- 500U --.

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

Page 4 of 7

PATENT NO. : 5,547,933

DATED : August 20, 1996

INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

- . column 11, line 41, "DN" should be -- DNA --.
- . column 14, line 10, "17 mer" should be -- 17-mer --.
- column 16, line 5, "128-mer" should be -- 128 20-mer --.
- column 16, line 46, "25" should be -- 25% -- .
- column 16, line 58, "<sup>125</sup>I labelled" should be -- <sup>125</sup>I-labelled --.
- column 16, line 58, after "added," insert -- and --.
- . column 17, line 40, after "procedures;" insert -- (4) --.
- column 17, line 62, "EOTA" should be -- EDTA --.
- column 18, line 16, "micomoles" should be -- picomoles --.
- column 18, line 33, "EgoRI" should be -- EcoRI --.
- column 18, line 44, "EcoRI/HindIII" should be -- EcoRI/HindIII --.
- column 20, line 7, "(NACl," should be -- (NaCl, --.
- column 20, line 17, "EPY" should be -- EPV --.
- column 21, line 67, coRI should be -- EcoRI --.
- column 22, line 19, "vector s" should be -- vector's --.
- column 22, line 49, "suu" should be -- supra --.

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

Page 5 of 7

PATENT NO. : 5,547,933  
DATED : August 20, 1996  
INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

column 23, line 43, "Sall" should be -- Sall --.

column 23, line 46, "Sall" should be -- Sall --.

column 24, line 51, "H and L" should be -- (H and L) --.

column 24, line 62, "he" should be -- the --.

column 26, line 6, "pDSVL-kEPO" should be -- pDSVL-MkEPO --.

column 26, line 8, "pDSVL-kEPO" should be -- pDSVL-MkEPO --.

column 26, line 14, "pOSVL" should be -- pDSVL --.

column 26, line 22, "(OHFR)" should be --(DHFR) --.

column 26, line 29, "NTX" should be -- MTX --.

column 26, line 30, "OHFR" should be -- DHFR --.

column 26, line 31, "OHFR" should be -- DHFR --.

column 26, line 32, "NTX" should be -- MTX --.

column 26, line 35, after "OHFR" should be -- DHFR --.

column 26, line 35, after "OHFR" should be -- DHFR --.



UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

Page 6 of 7

PATENT NO. : 5,547,933  
DATED : August 20, 1996  
INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

column 26, line 39, "pOSVL-NkE" should be --"pDSVL-MkE --.

column 26, line 39, "NTX" should be -- MTX --.

column 26, line 44, "NTX" should be -- MTX --.

column 26, line 45 "nN" should be -- nM --.

column 27, line 18, "pDSYL-gHuEPO" should be -- pDSVL-gHuEPO --.

column 27, line 30, "lie" should be -- line --.

column 27, line 32, "use" should be -- used --.

column 27, line 34, "replace" should be -- replaced --.

column 27, line 36, "m" should be -- mM --.

column 28, line 6, "pDSVL-GHuEPO" should be pDSVL-gHuEPO --.

column 28, line 7, "TX" should be -- MTX --.

column 29, line 56, "KEqI" should be -- KpnI --.

column 31, line 9, "Xhol" should be -- XhoI --.

column 31, line 26, "indIII" should be -- HindIII --.

column 31, line 26, "KpnI/BqIII" should be -- KpnI/BqII --.

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

Page 7 of 7

PATENT NO. : 5,547,933  
DATED : August 20, 1996  
INVENTOR(S) : Fu-Kuen Lin

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

column 31, line 53, "DYE" should be -- pYE --.

column 32, line 47, after "genotype" but before "pep4-3" insert -- & --.

column 32, line 55, "C34" should be -- (34 --.

column 33, line 60, "7000 U) g/kg" should be -- 7000U)  $\mu$ g/kg --.

column 34, line 55, "an" should be -- and --.

column 35, line 21, "interfetch" should be -- interferon --.

column 36, line 19, after "such as" insert -- in EPO assays or EPO antagonism. Antagonists of erythropoietin --.

column 37, line 10, "DA" should be -- DNA --.

column 37, line 10, second occurrence of "DA" should be -- DNA --.

column 37, line 50, "P.aeruginosa" should be -- P.aeruginosa --.

in the Claims, column 39, line 2, after "amount" insert -- of --.

Signed and Sealed this

Twenty-fourth Day of March, 1998

Attest:



BRUCE LEHMAN

Attesting Officer

Commissioner of Patents and Trademarks

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 5,547,933  
APPLICATION NO. : 08/487774  
DATED : August 20, 1996  
INVENTOR(S) : Fu-Kuen Lin


Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Claim 14, column 40, line 9, "product" should be --patient--.


Signed and Sealed this

Seventeenth Day of April, 2007

A handwritten signature in black ink on a light gray dotted background. The signature reads "Jon W. Dudas" in a cursive style.

JON W. DUDAS

*Director of the United States Patent and Trademark Office*



**THE UNITED STATES OF AMERICA**

**TO ALL TO WHOM THESE PRESENTS SHALL COME:**

**UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office**

November 02, 1998

THIS IS TO CERTIFY THAT ANNEXED IS A TRUE COPY FROM THE RECORDS  
OF THIS OFFICE OF THE FILE WRAPPER AND CONTENTS OF:

APPLICATION NUMBER: 08/487,774

ISSUING DATE: June 07, 1995

PATENT NUMBER: 5,547,933

ISSUANCE DATE: August 20, 1996

TITLE OF INVENTION:

FUNCTION OF ERYTHROPOIETIN

(S):

By Authority of the  
**COMMISSIONER OF PATENTS AND TRADEMARKS**

**N. WOODSON**

**Certifying Officer**