



Mercury in canned tuna: white versus light and temporal variation

Joanna Burger^{a,b,*} and Michael Gochfeld^{b,c}

^a *Nelson Biological Laboratory, Division of Life Sciences, Rutgers University, 604 Allison Road, Nelson Hall, Piscataway, NJ 08854-8082, USA*

^b *Environmental and Occupational Health Sciences, Institute and Consortium for Risk Evaluation with Stakeholder Participation, Piscataway, NJ 08854, USA*

^c *UMDNJ–Robert Wood Johnson Medical School, Environmental and Community Medicine, Piscataway, NJ 08854, USA*

Received 17 August 2003; received in revised form 26 November 2003; accepted 3 December 2003

Abstract

There are abundant data and advisories for mercury levels in wild fish, but far fewer for commercial fish that compose a large majority of the fish most people eat. Until recently, relatively little attention has been devoted to examining mercury in canned tuna, despite its great importance in human diets. There is substantial media coverage of the benefits and risk from fish consumption, but few peer-reviewed data on canned tuna, the most commonly consumed fish in the United States. In this paper, we examine the levels of total mercury in canned tuna obtained from a New Jersey grocery store from 1998 to 2003, looking for temporal consistency within this data set and particularly for comparison with the Food and Drug Administration's 1991 study. We analyzed 168 cans individually for total mercury. All values are reported as parts per million (= µg/g) on a wet weight basis. In a subset of samples analyzed for total and inorganic mercury, the inorganic mercury was below detection levels; hence at least 89% of the mercury can be considered methylmercury. We found that white-style tuna had significantly more total mercury (mean 0.407 ppm) than light-style tuna (mean 0.118 ppm), presumably reflecting that "white" tuna is albacore, a species relatively larger than the skipjack tuna, which is commonly available as "light" or "chunk light." The maximum mercury in a can was 0.997 ppm, but 25% of white tuna samples exceeded 0.5 ppm. Data suggest a slight increase in levels since 1991, and mercury levels were significantly higher in 2001 than in other years. The mean level of mercury in white tuna (mean 0.407 ppm) was significantly higher than the mean value of 0.17 ppm currently used by the U.S. Food and Drug Administration (FDA) in its risk assessment and public information. There were no significant differences in mercury levels in tuna packed in oil compared to water. Draining contents had no effect on mercury levels, and the fluid, both oil and water, contained little mercury. These data indicate that people who eat canned tuna frequently can choose light tuna and reduce their mercury intake. Canned mackerel had much lower levels of mercury than tuna. Since cans of white tuna frequently exceed the FDA's original action level of 0.5 ppm, it would be prudent to continue some systematic monitoring of the nation's canned fish supply, particularly as the targets of commercial fisheries inevitably change as certain stocks become depleted.

© 2003 Elsevier Inc. All rights reserved.

Keywords: Methylmercury; Mercury; Canned tuna; Exposure; Risk; Food and drug administration; Consumption advisories; Albacore

1. Introduction

Fish provide a healthful source of dietary protein, and are relatively low in cholesterol and high in omega-3 (n-3) fatty acids (National Research Council, 2000). Several studies have documented the long-term cardio-protective benefits for adults as well as the reproductive benefits of eating fish. However, benefits may be offset by the presence of contaminants, particularly methylmercury (MeHg) and polychlorinated biphenyls (PCBs),

which have been reported in many fish species from many localities (Environmental Protection Agency [EPA], 1998). Over more than three decades studies have shown positive correlations between mercury levels in humans and fish consumption (Bjornberg et al., 2003). A recent review of data from the National Health and Nutrition Examination Survey (NHANES) found that women who ate three or more servings of fish a month had a mercury level four-fold higher than that in the reference population (Schober et al., 2003).

Contaminants in fish are particularly detrimental for developing fetuses and young children (Jacobson et al., 1989, 1990; Institute of Medicine [IOM], 1991; Sparks

*Corresponding author. Fax: 732-445-5870.

E-mail address: burger@biology.rutgers.edu (J. Burger).

and Shepherd, 1994; Jacobson and Jacobson, 1996; Schantz, 1996; National Research Council, 2000). Methylmercury interferes with the architecture of the developing brain, disrupting microtubule assembly (Graff et al., 1993) and interfering with the temporal sequencing of cell adhesion molecules that guide neuronal migration and connections (Dey et al., 1999). Some studies have shown an association between contaminant levels in fish, fish consumption by pregnant women, and deficits in neurobehavioral development in children (Weihe et al., 1996; Jacobson and Jacobson, 1996; Grandjean et al., 1998).

1.1. *Advisories*

In many states contaminant levels in some recreationally caught freshwater fish are sufficiently high to cause adverse health effects (IOM, 1991), which has led to managing risk by issuing consumption advisories for many water bodies and fish species. Fish advisories stimulated our studies of anglers to determine consumption patterns, perceptions of risk by anglers, and their compliance with advisories (Burger et al., 1993, 1999a, b). Despite the abundance of advisories aimed at fishermen, there is a paucity of advice for consumers of commercial fish, although this makes up about 95% of all consumption in the United States (Stern et al., 1996).

In 2001, the U.S. Food and Drug Administration (FDA) issued a consumption advisory, based on mercury, that pregnant women and women of child-bearing age who may become pregnant should avoid eating four species (or species groups) of saltwater fish: shark, swordfish, king mackerel, and tilefish (FDA, 2001a). In 2002 the FDA's advisory committee recommended expanding the advice to specifically mention tuna, but that has not been done to date. The FDA is currently (November and December 2003) reviewing its advisory.

Methylmercury is among the best known of environmental contaminants, engendering an extensive literature (Mahaffey, 1999; National Research Council, 2000). The risks from mercury in fish gained widespread media coverage in 2003 (e.g., Raines, 2003; Gorman, 2003), calling attention to canned tuna as a source of excessive mercury.

The present study was begun to assist New Jersey's Mercury Task Force in developing recommendations about advisories and monitoring (NJ Task Force, 2002). It also complemented our studies of wild-caught fish in New York (Burger et al., 1993), New Jersey (Burger et al., 1999a), Puerto Rico (Burger and Gochfeld, 1991), and South Carolina (Burger et al., 1999b). In this paper, we examine the levels of mercury in national brands of canned tuna purchased from a New Jersey grocery store chain from 1998 to 2003. This paper addresses the levels

of mercury in different styles (solid white, chunk white, and chunk light tuna) and examines interyear variation, including comparison with the FDA's 1991 study (Yess, 1993). We also tested differences between oil- and water-packed tuna and between drained and undrained contents.

1.2. *Benefits of fish consumption*

The question of risk from eating fish is complicated, however, by the positive health and social benefits of consuming fish (Toth and Brown, 1997). Fish provide omega-3 (n-3) fatty acids, which reduce cholesterol levels (Hunter et al., 1988; Kimbrough, 1991; Horn, 1992; Anderson and Wiener, 1995). For some people fish may be their main source of protein, while for others, it may be the healthiest source. Fish consumption has been associated with improved pregnancy outcomes, including fewer preterm and low-birth-weight deliveries (Olsen and Secher, 2002), attributable in part, at least, to n-3 fatty acids (Allen and Harris, 2001). On the other hand Guallar et al. (2002) indicated that mercury might offset the cardioprotective benefits for adults. Thus, assessing the levels of mercury in commercial fish is important from a public health perspective.

2. *Methods*

2.1. *Source and types of canned tuna samples*

Canned tuna was purchased from one central New Jersey chain grocery store. Our sampling regime varied over time, and involved purchasing three to five cans each of several different national brands of tuna (in water) each year from 1998 to 2003. After initial analyses revealed higher mercury levels in white than in light tuna, in later years we oversampled chunk white and solid white. Sample sizes and styles are given in Table 1. Mercury was analyzed in the Environmental and Occupational Health Sciences Institute Elemental Analysis Laboratory. In several years we examined mercury levels in tuna packed in oil and compared levels in drained and undrained samples. In 2001 we also compared mercury levels in drained and undrained canned mackerel ($n = 9$).

We analyzed tuna from 168 cans including 86 solid white, 37 chunk white, and 45 chunk light. These designations are standardized by the Food and Drug Administration (FDA, 1977) with reference to a Munsell gradient of darkness. White "is limited to the species *Thunnus germon* (albacore), and is not darker than Munsell value 6.3". Light tuna ranges from Munsell 6.3 to not darker than 5.3. Solid consists of "loins cut in transverse segments to which no free fragments are

Table 1
Total mercury in canned tuna (1998–2003)

	N	Median	Mean	SD	SE	CV	Max
Entire sample	168	0.311	0.330	0.199	0.015	60.3	0.997
All white tuna	123	0.368	0.407	0.167	0.015	41.0	0.997
Chunk white	37	0.315	0.355	0.166	0.027	46.8	0.997
Solid white	86	0.4	0.429	0.164	0.018	38.2	0.783
All light tuna ^a	45	0.087	0.118	0.099	0.015	83.9	0.447
Comparing all white with all light					$\chi^2 = 78.9^b$	$P < 0.0001$	
Comparing all chunk with all solid white					$\chi^2 = 6.39$	$P = 0.010$	
All tuna in water	136		0.346	0.196	0.017	56.7	0.997
All tuna in oil	32		0.259	0.197	0.035	76.0	0.719
Comparing all water with all oil					$\chi^2 = 5.54$	$P = 0.019$	
All light in water ^d	26	0.072	0.096	0.062	0.012	64.7	0.244
All white in water	110	0.368	0.405	0.168	0.016	41.5	0.997
Chunk white in water	37	0.315	0.355	0.166	0.027	46.6	0.997
Solid white in water	73	0.401	0.431	0.165	0.019	38.3	0.783
Comparing light with white in water					$\chi^2 = 58.5$	$P < 0.0001$	
Comparing chunk versus solid white in water					$\chi^2 = 6.3$	$P = 0.012$	
Oil							
Chunk light ^d	19	0.112	0.149	0.130	0.030	87.5	0.447
Solid white	13	0.369	0.419	0.166	0.046	39.6	0.719
Comparing light with white in oil					$\chi^2 = 15.2$	$P < 0.0001$	

All values are in ppm ($\mu\text{g/g}$) on wet weight basis.

^aAll of the light tuna was designated "chunk light."

^bKruskal–Wallis non parametric one-way analysis of variance computed using the Wilcoxon Option in Proc NPARIWAY of SAS, yielding a χ^2 value.

Table 2
Interlaboratory comparisons on split samples of canned tuna (New Jersey)

Year	N	EOHSI laboratory	Lancaster laboratory ^a	Quebec laboratory ^b	Correlations with		Percent difference of means ^c	
		Mean \pm SE	Mean \pm SE	Mean \pm SE	Lancaster	Quebec	Lancaster	Quebec
2001	8 ^d	0.701 \pm 0.03	0.691 \pm 0.06	0.663 \pm 0.05	0.95	0.72	4.1%	5.4%
2002	20	0.239 \pm 0.03	No longer providing service	0.265 \pm 0.03		0.93		9.8%
2003	18	0.301 \pm 0.06	No longer providing service	0.286 \pm 0.05		0.93		6.6%

The samples sent out in 2001 were all high in mercury (> 0.5 ppm). The 2002 and 2003 were randomly chosen split samples.

^aLancaster's Food Service Laboratory ceased providing this service after 2001.

^bInstitut national de santé publique du Québec.

^cPercent difference between the mean values in the outside reference laboratory and the EOHSI mean value.

^dEight samples all over 0.50 ppm were submitted for confirmation in 2001.

added" while chunk consists of "a mixture of pieces in which the original muscle structure is retained" (FDA, 1977). Light tuna comprises mainly skipjack (*Katsuwonus pelamis*), which is generally a smaller species than albacore (which is now generally considered *Thunnus alahunga*).

2.2. Analysis

Sample preparation, digestion, and analysis were done in three separate rooms to minimize cross

contamination. All analyses were performed in the Elemental Analysis Laboratory of the Environmental and Occupational Health Sciences Institute, which is a partnership of Rutgers University and Robert Wood Johnson Medical School. A subset of samples were analyzed at outside reference laboratories for quality assurance (see Table 2). All analyses were done on individual cans; compositing was not performed. Contents were drained and samples were weighed to 0.1 mg. All analyses and calculations were carried out to four significant figures. All laboratory equipment and

containers were washed in 10% HNO₃ solution prior to each use and rinsed with deionized water. Fish was digested in Ultrex ultrapure nitric acid in a microwave (MD 2000 CEM), using a digestion protocol of three stages of 10 min each under 50, 100, and 150 pounds per square inch (3.5, 7, and 10.6 kg/cm²) at 70 × power. Digested samples were subsequently diluted in 100 ml deionized water. Mercury was analyzed by cold vapor atomic absorption spectrometry using the HGS-4 analyzer. In 2002 and 2003 analyses were also performed using a Lumex atomic absorption spectrometer with Zeeman background correction, which provided better precision near the detection level. Correlations between the HGA-4 and Lumex exceeded 0.92 and mean values of replicates were within 3%.

The method detection limit (MDL) ranged from 0.05 ppm with the HGA-4 to 0.03 with the Lumex and the few nondetectable values were set at half the MDL. For water drained from cans, the MDL was 0.001 ppm. We used Kruskal–Wallis nonparametric one-way analysis of variance (SAS-NPAR1WAY with Wilcoxon option) to examine for differences among independent variables (Statistical Analysis System [SAS], 1995) and nonparametric Kendall tau procedure for testing correlations. The level for significance was designated as <0.05. All results are reported as parts per million (ppm), equivalent to micrograms per gram on a wet weight basis, rounded to three significant figures.

2.3. Quality control analyses

All specimens were run in batches that included blanks, a standard calibration curve, and spiked specimens. The accepted recoveries for spikes ranged from 85% to 115%; batches outside these limits were rerun. The coefficient of variation (C.V.) on replicate samples ranged from 2% to 12%. Further quality control included periodic blind analysis of an aliquot from a large sample of known concentrations and blind runs of replicate samples. Analyses from earlier years were rerun and confirmed with batches in later years. Replicates were run for half of the cans, and all digests were analyzed in duplicate with mean values used. We also measured the moisture content.

We used two external laboratories to analyze split samples. In 2001 samples were submitted to the Lancaster Laboratories, Inc., Food Service Division (which closed in 2002) and to the Institut National de Santé Publique du Québec (Quebec) for quality control (Table 2). In 2002 and 2003 samples were also sent to the Quebec laboratory, which runs the international proficiency testing program for mercury. High results (>0.50 ppm) from early years were confirmed in the outside laboratories.

3. Results

3.1. Quality control

Quality control is an essential part of any laboratory quality assurance program. The quality control samples (blanks, standards) were within specifications, and the replicate data used were all within 15%. However, we were surprised by the number of replicates that exceeded 10%. This prompted us to test several cans repeatedly, demonstrating that the ratio of maximum to minimum in different parts of a single can vary by more than 10%. The average difference between duplicates (on the same extract) was 0.2%. The average difference between replicates (separate digests of same sample) was 0.78%.

The comparison between the EOHSI and the outside laboratories shows generally very close agreement across laboratories, despite differences in method (Table 2). The Lancaster results were obtained on dried specimens corrected for moisture content. The Quebec results were reported to the nearest 0.05 ppm with an MDL of 0.05. Correlations were generally high and means were in close agreement.

3.2. Total mercury

The main findings of our study were the consistently higher levels in white tuna (overall mean 0.407 ppm; median 0.368 ppm) than in light tuna (mean 0.118 ppm; median 0.087 ppm; Table 1). We found a somewhat higher mean (0.429 ppm) for solid white than for chunk white (0.355 ppm), but this varied across years. The level of 0.3 ppm was exceeded by 67% of the white and 7% of the light tuna samples; 25% of the white, but none of the light exceeded 0.5 ppm. The maximum value for total mercury in our sample was 0.997 ppm. We found significant variation among years; samples purchased in 2001 were higher than in earlier or later years of the study (Table 3). There were no differences in the concentrations of mercury for tuna packed in oil or in

Table 3
Mercury in canned tuna: comparison among years for white tuna in water

Year	N	Mean
1998	6	0.319
1999	21	0.314
2000	18	0.212
2001	27	0.517
2002	12	0.332
2003	26	0.475
Total N	110	Grand mean = 0.361
Among years (1998–2003)		
Kruskal–Wallis χ^2	33.6	
Probability	$P < 0.0001$	

All values are for total mercury in ppm ($\mu\text{g/g}$) on wet weight basis.

water, and no differences for whether it was drained or undrained (Table 4). Similarly, there were no differences in the mercury concentrations of drained versus undrained mackerel, another readily available canned fish (Table 4), which had levels almost an order of magnitude lower than in white tuna.

Because we oversampled white tuna (both solid and chunk) in water, we do not emphasize a single mean value for all canned tuna. Multiple regression analysis for total mercury in water-packed tuna was tested with the general linear models procedure (PROC GLM) of SAS, using brand, style (white versus light), and year as independent class variables. The overall model was significant ($F = 13.2$, $P < 0.001$; Table 5). Both style ($F = 84$, $P < 0.0001$) and year ($F = 7.02$, $P < 0.0001$) were significant contributors to the overall r^2 of 0.6, while brand did not enter significantly ($F = 1.7$, $P > 0.20$). The sample size for oil-packed was too small for this analysis.

3.3. Yearly variation

No attempt was made to achieve a balanced design of style over years. Data broken down by years is presented

Table 4
Comparison of mercury levels in canned tuna and mackerel before and after draining

	Number	Mean \pm SE	Kruskal–Wallis χ^2	(<i>P</i> value)
Canned tuna				
Undrained	11	0.43 \pm 0.03	0.8	NS
Drained	11	0.46 \pm 0.15		
Canned mackerel				
Undrained	9	0.033 \pm 0.003	1.1	NS
Drained	9	0.026 \pm 0.004		

All values are for total mercury in ppm ($\mu\text{g/g}$) on wet weight basis.

Table 5
Comparison of current results with the FDA's 1991 study (Yess, 1993)

		Chunk light	Chunk white	Solid white	All white	Water ^a	Oil ^a
1991 FDA study (Yess, 1993)							
Methylmercury (analyzed)	Mean	0.100	0.310	0.260	0.271	0.191	0.079
	SD	0.110	0.170	0.160	0.160	0.160	0.100
Current study							
Methylmercury (calculated based on 89%) ^b	Mean	0.132	0.353	0.480 ^c	0.412	0.354	0.245
Methylmercury (calculated) excluding year 2001	Mean	0.132	0.375	0.442 ^c	0.418	0.352	0.242
Analyzed methylmercury for 2003 only ^d	Mean	0.040	0.450	0.329	0.355	0.243	0.233

All values are in ppm ($\mu\text{g/g}$) on wet weight basis.

^aOil and water results should not be compared because of differential sampling of white and light in different years. Our sample was only 27% light compared to 54% for Yess (1993).

^bOur results estimate 89% of mercury as methylmercury, agreeing closely with other estimates. Analyzed results were divided by 1.12 to estimate the MeHg.

^cIndicates that results from this study are more than 1 SD greater than 1991 results (Yess, 1993).

^dAnalyzed by the Institut national de sante publique due Quebec.

in Table 3. The highest single category for any year was the mean value of 0.540 for chunk white in water in 2003. The lowest was for chunk light in oil of 0.074 in 2000. Our mean values for light and chunk white were similar to the 1991 data, but our mean for solid white was significantly higher ($P < 0.05$). We also compared our methylmercury results, both calculated and analyzed, with the FDA's 1991 data (Table 5), finding that our results for solid white tuna (1998–2003) were significantly higher than the 1991 results (Yess, 1993). Our results for chunk light and chunk white were slightly, but not significantly higher.

3.4. Oil versus water

Many brands of tuna are packed in both oil and water. Oil is usually vegetable oil (sometimes olive oil). We compared each of the three styles packed in oil versus water (Table 1). None of the comparisons were significant ($P > 0.10$), although oil-packed tuna averaged slightly lower levels for white and higher levels for light ($P > 0.10$).

3.5. Drained versus undrained

We drained all cans prior to removing tissue for analysis. In a subset of cans we compared drained and undrained results. Draining had no consistent effect on mercury content (Table 4). Likewise, the liquid component, when analyzed, had very low levels of mercury, usually below the MDL for liquid of 0.001 ppm with a maximum of 0.07 ppm.

3.6. Methylmercury content

Forty samples were analyzed by the Quebec laboratory for total and inorganic mercury, calculating

methylmercury by subtraction. The inorganic component was below the detection level of 0.05 ppm for all samples. Using the convention of half the detection level (i.e., 0.025) for the inorganic component yields an estimate of the organic mercury content of 89% for samples, where total mercury was 0.10 ppm or greater. We show the calculated MeHg values for 6 years and 5 years (excluding 2001), as well as the actual analyzed values for 2003 (see Table 5).

3.7. Provenance

Commercial tuna sold in the United States is caught mainly in the Pacific. Most cans of white tuna specified “albacore.” Only about $\frac{1}{4}$ of the cans indicate a country of origin—usually Thailand, although a few mentioned New Guinea or the Philippines. Other companies obtain tuna from American Samoa and Ecuador. The fish may have been caught near these countries or may simply have been landed or canned there. There were similar batch numbers on cans from Thailand representing more than one brand, indicating that canneries service more than one company. In most cases, however, we have no idea where the fish were caught other than in the Pacific, or whether the source of tuna shifted from year to year, which might partly explain annual variation.

4. Discussion

Canned tuna amounts to about 20–25% of all seafood consumed in the United States, with average per capita consumption in 2000–2001 of about 1.5 kg (National Marine Fisheries Service [NMFS], 1989; National Oceanic Atmospheric Administration [NOAA], 2002). Although most Americans have relatively low exposure, a small percentage, accounting ultimately for several million people, have excessive mercury levels due in part to fish consumption including canned tuna (Stern et al., 2001; Hightower and Moore, 2002). Our results are consistent with previously published results regarding differences among tuna styles, and although the average size of tuna may have declined, the mercury levels in canned tuna have not, and may have increased. Although our sample size is miniscule compared to the number of cans of tuna sold in the United States, it is relatively large compared with most published studies.

4.1. Mercury in tuna

Many studies have shown that mercury is bioamplified in the food chain with high-trophic-level predatory species (such as shark, swordfish, and tuna) having higher mercury (and organochlorines) than herbivorous species (Braune, 1987; Lacerda et al., 1994; Park and

Curtis, 1997; Burger et al., 2001). Within a species, older and larger fish also have higher levels, as Boush and Thieleke (1983) showed for tuna.

Although there are several published reports of mercury in canned tuna, most studies have not distinguished between different styles of tuna; hence comparisons with the literature are difficult. It is often assumed (even in fish advisories) that fresh tuna or tuna steaks have higher mercury levels than canned tuna, but the literature does not provide clear evidence for this. The FDA Web site (FDA, 2001b) reports a value for methylmercury of 0.32 ppm for fresh or frozen tuna compared with its value of 0.17 ppm for canned tuna. However, mean values for white style canned tuna exceed 0.3 (Yess, 1993; this paper).

The National Marine Fisheries Service (1978) measured total mercury in 250 individuals of several species of tuna, yielding a median of 0.12 ppm and a maximum of 0.87 ppm. However, Plessi et al. (2001) reported an average of 0.249 in bluefin tuna (*Thunnus thynnus*) purchased from a fish wholesaler in Italy. Nakagawa et al. (1997) reported levels of mercury in fresh tuna from Japan of 1.11 ppm. Storelli et al. (2002) found levels of 0.85–1.45 ppm (wet weight) in albacore from the Mediterranean (average 1.17 ppm) with 78% exceeding 1 ppm. Consumer Reports (2001) recently published an average value of 0.25 ppm methylmercury in 24 samples of fresh tuna. Moreover, there is evidence of variation among oceans and countries, probably related to the source of the fishery, species, and fish size.

4.2. Mercury in canned tuna

The FDA reports that methylmercury levels in canned tuna fish are normally in the range 0.1–0.2 ppm (Carrington et al., 1997), and the value of 0.17 is given on its Web site as the average for canned tuna (FDA, 2001b). This is based on the study of Yess (1993), which bears closer examination (next section).

Most studies report only total mercury. Yallouz et al. (2001) analyzed data from 39 cans of tuna representing five brands from Brazil, without details on style. The grand mean was 0.65 ppm, with 51% of cans exceeding 0.5 ppm and 15% greater than 1 ppm. Voegborlo et al. (1999) analyzed 50 cans of chunk tuna from the Mediterranean and reported a mean of 0.29 ppm (SD = 0.12), based only on the 20 samples that exceeded the MDL of 0.2. Hence the true mean for their sample would have been substantially lower; we calculated it as 0.12 ppm setting the 30 nondetected values to half the MDL. This is in close agreement with both our data and those of Yess (1993) for light tuna. Acra et al. (1981) reported a mean of 0.30 $\mu\text{g/g}$ (range 0.25–0.49) for canned tuna from Lebanon. Other results from this period include mean values of 0.3 ppm (Kamps et al., 1972; Parvaneh, 1979), but the style was not specified.

White tuna is albacore, a species somewhat larger than those that are canned as light tuna. Albacore generally weigh 10–20 kg, but can reach 40 kg. Skipjack, the main component of light tuna, accounts for about half of all tuna harvested worldwide. It averages about 8–10 kg in weight, occasionally reaching 30 kg (International Foundation for Conservation of Natural Resources [IFCNR], 2003). Albacore tends to have high mercury levels (Storelli et al., 2002).

4.3. FDA's 1991 study

Between 1978 and 1990, FDA determined methylmercury in 42 samples of canned tuna (ranging from <0.10 to 0.67 ppm, with an average of 0.14 ppm reported in Yess, 1993). The FDA's 1991 study (Yess, 1993) is the single largest and best study of mercury in canned tuna. Yess (1993) analyzed 220 samples, each a composite of 12 cans (12 additional cans were archived). The cans were purchased in 1991 from around the United States. The sampling was nonrandom and emphasized water-packed tuna and popular brands. She found that samples ranged from the detection level of 0.1 ppm up to 0.75 ppm of methylmercury; 3.6% of her composite samples exceeded 0.5 ppm methylmercury. The overall mean was reported as 0.17 ppm (median of 0.15 and a maximum of 0.90 ppm; Yess, 1993). Yess (1993) also reported that chunk white tuna averaged 0.31 (SD=0.17) and solid white averaged 0.26 (SD 0.16), while chunk light tuna and chunk averaged 0.10 (SD=0.11). The light styles made up 2/3 of her sample, resulting in an unweighted mean of 0.17. She arbitrarily set all nondetectable values to 0 rather than to the more commonly used convention of one-half the MDL or 0.05 ppm; hence her results are biased downward. None of the samples exceeded the current 1 ppm FDA action level (Yess, 1993).

Based on the Yess (1993) paper, the FDA posts 0.17 ppm on its Web site as the value for canned tuna (FDA, 2001b) and uses this value in its exposure and risk assessments (Carrington et al., 1997; Carrington and Bolger, 2002; Bolger and Schwetz, 2002). No mention is made of the white versus light discrepancy, even though it was recognized by Yess (1993).

4.4. Oil versus water packing

We did not find a difference in the mercury concentration of oil-packed versus water-packed tuna within each of the three styles (Table 1). Yess (1993) reported a mean of 0.06 ppm for 26 oil-packed composites versus 0.11 ppm for 106 water-packed composites of chunk light tuna. However, the FDA study blended the oil into the fish sample, whereas water was drained. Hence this would serve to dilute the mercury somewhat.

4.5. Drained versus undrained

Advice to consumers sometimes mentions draining to reduce contaminants. In 2001 we analyzed 11 cans of tuna undrained and again after draining. The results (Table 4) show no appreciable change ($P>0.25$). All other analyses were performed after the contents were manually drained, but not squeezed. Liquid volume varied among cans.

4.6. Interyear variation

We found a statistically significant peak in the 2001 samples of white tuna ($\chi^2 = 33$; $P<0.001$). This may be a statistical artifact of sample size, but more likely reflects exploitation of a different population of tuna due to the vagaries of the fish movements and fishery targets.

For several reasons, when we began the study, we expected a decline in mercury levels over time. We anticipated that a decline in mercury contamination through the gradual reduction in mercury use and industrial effluents would be reflected in a decline in the mercury content of tuna. We suspected also that mercury levels might have declined as a result of overfishing large tuna and the need to target smaller fish (Safina, 1998). Moreover, in New Jersey common terns (*Sterna hirundo*), a fish-eating seabird, we found that mercury levels in eggs had declined since the early 1970s (Burger and Gochfeld, in press).

In recent years, however, mercury emissions, mainly from coal-fired power plants have actually increased (NJDEP, 2001), resulting in short- and long-range atmospheric transport, which may offset declines in use and effluent. To the extent our data show a temporal trend, it has been upward since 1991 (Burger and Gochfeld, in press). Yess (1993) suggested that FDA data might have shown a decline from a calculated mean of 0.21 for methylmercury in 1973 to 0.17 in 1991. However, the composition of the 1973 sample with respect to white and light was not given.

4.7. Methylmercury in tuna

This study was not primarily designed to evaluate methylmercury per se, since we initially followed the conventional wisdom that about 90% of the mercury in fish is methylmercury. However, we did measure the methylmercury content in a subset of cans and found that methylmercury composed 83–89%. Considering only cans with total mercury >0.10, the value was about 89%. Storelli et al. (2002) reported a range of 75–100% MeHg (average 91%) and Hall (1974) reported a mean of 89%. Using 89% results in a correction factor of 1.12, and the methylmercury content of our samples can be estimated by dividing total mercury by 1.12. It is not

clear how much of the methylmercury in tuna (or other marine fish) is anthropogenic in origin.

4.8. Action levels and tuna as an exemption

The regulatory control of tuna, as well as shark and swordfish, is problematic. For example, in Japan, Canada, and Europe, tuna are exempted from the 0.4 ppm or 0.5 ppm regulations (Nakagawa et al., 1997; Canadian Food Inspection Agency, 2002; Plessi et al., 2001). In the U.S., the FDA (1987) level of 1 ppm is normally high enough so that tuna falls below this level, although this was not always the case. In 1969 FDA developed administrative guidance of 0.5 ppm for commercial fish, and in the early 1970's millions of cans of fish exceeding this level were seized. In 1974 this was converted to an Action Level, but in 1979 FDA raised its Action Level to 1.0 ppm.

Many countries have set the maximum permitted level of mercury in fish at 0.5 ppm, including Australia (Denton and Burdon-Jones, 1996), Canada (Institute of Medicine [IOM], 1991), Sweden (Hylander et al., 1994), and the United Kingdom (Collings et al., 1996). Many states in the U.S. have also set 0.5 ppm or lower limits, including Florida (Lange et al., 1994), Maine (DiFranco and Mower, 1994), Minnesota (Minnesota Department of Health [MDH], 1997), and Wisconsin (Gerstenberger et al., 1993). Of the cans of tuna we examined, 50% exceeded 0.3 ppm and 25% of white style exceeded 0.5 ppm. In 2001, half of the samples were above 0.50 ppm, the limit many states and countries set for safe consumption.

People must balance potential mercury exposure with the need for a healthy diet. Our study provides additional data that the levels of mercury in tuna can be both variable and high, lessening the predictability of the mercury a pregnant woman consumes when eating canned tuna.

The variability in mercury levels in canned tuna is particularly worrisome because there are sensitive periods for neurobehavioral development when the developing fetus is more susceptible to the disruptive effects of mercury, and when a single pulsed dose may have more of an effect than the same amount over a longer timeframe (Dey et al., 1999).

4.9. Risk and risk reduction

Recently, the Agency for Toxic Substances and Disease Registry [ATSDR] (1999) has proposed a Minimal Risk Level of 0.3 $\mu\text{g}/\text{kg}/\text{day}$ mercury based on the Seychelles neurodevelopmental study (Davidson et al., 1998), without incorporating an uncertainty factor for interindividual variation. That study has not found neurobehavioral deficits in a fish-eating population (Myers et al., 2003). The Faroe Island study (Weihe

et al., 1996; Grandjean et al., 1998) supports a lower Reference Dose (RfD), which the EPA has calculated at 0.1 $\mu\text{g}/\text{kg}/\text{day}$, and Stern (1993) computed an RfD of 0.07 based on data from Iraq. The National Research Council (2000) concluded that 0.1 $\mu\text{g}/\text{kg}/\text{day}$ was scientifically sound.

There is often a gap between how the fish-consuming public and the agencies responsible for advisories perceive risk (Reinert et al., 1991; Ebert, 1996; Burger, 2000). The public views eating fish as a less serious hazard than does the scientist or environmental manager, but more seriously than some regulators. People continue to consume the fish even though they know about the consumption advisories or bans (Reinert et al., 1991; Burger and Gochfeld, 1991; Burger et al., 1992, 1999a; Velicer and Knuth, 1994; May and Burger, 1996; Burger et al., 2001). Until 2003, however, almost no attention was devoted in the media to the risks from tuna fish.

Reducing risk from the consumption of fish is obviously a partnership between risk assessors and resource managers (Ebert, 1996), as well as the public (Burger, 1998), and involves balancing the risks versus the advantages (Egeland and Middaugh, 1997; Bolger and Schwetz, 2002). We suggest that the problem of risk perception concerning tuna will be particularly difficult because it is such a common and accepted food and has always been considered safe and healthy. Over the years the tuna industry has provided a significant reduction in risk by reducing the amount of tuna in cans by about 15%, to the current industry standard of 6 oz (170 g).

4.10. Recommendations

Our data indicate that people who are not in high-risk categories (most adults and adolescents) may consume several cans of light style tuna a week and white tuna occasionally with impunity. Pregnant or soon-to-be pregnant women, infants, and small children should limit canned tuna intake and select light rather than white tuna. These data (particularly their variability) indicate the desirability of instituting a regular monitoring program for commercial tuna and other species. Whereas canned fish is nutritious, tasty, and convenient, we found that mercury levels in canned mackerel were substantially lower than in tuna. FDA and other agencies should clarify the difference between mercury levels in white versus light tuna.

Acknowledgments

We thank T. Shukla and C. Dixon for mercury analysis and computer assistance, and A. Stern, K. K. Pflugh, B. Buckley, C. Chess, C. Powers, B. D. Goldstein, B. Friedlander, C. Powers, and A. Upton

for comments on the research and manuscript. Alain LeBlanc (Institut national de santé publique du Québec) provided valuable quality control assistance. Our research with fish consumption was also influenced by discussions with members of the Savannah River Site's (SRS) Citizen's Advisory Board. This research was partly funded by the Consortium for Risk Evaluation with Stakeholder Participation (CRESP) through the Department of Energy (AI DE-FC01-95EW55084, DE-FG 26-00NT 40938), NIESH (ESO 5022), and the Environmental and Occupational Health Sciences Institute. The results, conclusions, and interpretations reported herein are the sole responsibility of the authors, and should not in any way be interpreted as representing the views of the funding agencies.

References

- Acra, A., Namaan, S., Raffoul, Z., 1981. Total mercury levels in canned and frozen fish imported into Lebanon. *Bull. Environ. Contam. Toxicol.* 27, 209–212.
- Agency for Toxic Substances and Disease Registry [ATSDR], 1999. Toxicological Profile for Mercury. Centers for Disease Control, Atlanta, GA.
- Allen, K.G., Harris, M.A., 2001. The role of n-3 fatty acids in gestation and parturition. *Exp. Biol. Med.* 226, 498–506.
- Anderson, P.D., Wiener, J.B., 1995. Eating fish. In: Graham, J.D., Wiener, J.B. (Eds.), *Risk Versus Risk: Tradeoffs in Protecting Health and the Environment*. Harvard University Press, Cambridge, MA.
- Bjornberg, K.A., Vahter, M., Petersson-Grawe, K., Glynn, A., Cnattingius, S., Darnerud, P.O., Atuma, S., Aune, M., Becker, W., Berglund, M., 2003. Methyl mercury and inorganic mercury in Swedish pregnant women and in cord blood: influence of fish consumption. *Environ. Health Perspect.* 111, 637–641.
- Bolger, P.M., Schwetz, B.A., 2002. Mercury and health. *New Engl. J. Med.* 347, 1735–1736.
- Boush, G., Thieleke, J.R., 1983. Total mercury content in yellowfin and bigeye tuna. *Bull. Environ. Contam. Toxicol.* 39, 291–297.
- Braune, B.M., 1987. Mercury accumulation in relation to size and age of Atlantic herring (*Clupea harengus harengus*) from the south-western Bay of Fundy, Canada. *Arch. Environ. Contam. Toxicol.* 16, 311–320.
- Burger, J., 1998. Attitudes about recreation, environmental problems, and estuarine health along the New Jersey shore. *Environ. Manage.* 22, 869–876.
- Burger, J., 2000. Consumption advisories and compliance: the fishing public and the deamplification of risk. *J. Environ. Planning Manage.* 43, 471–488.
- Burger, J., Gochfeld, M., 1991. Fishing a superfund site: dissonance and risk perception of environmental hazards by fishermen in Puerto Rico. *Risk Anal.* 11, 269–277.
- Burger, J., Gochfeld, M., Metal levels in eggs of common terns (*Sterna hirundo*) in New Jersey: temporal trends from 1971 to 2002. *Environ. Research*, in press.
- Burger, J., Cooper, K., Gochfeld, M., 1992. Exposure assessment for heavy metal ingestion from a sport fish in Puerto Rico: estimating risk for local fishermen. *J. Toxicol. Environ. Health* 36, 355–365.
- Burger, J., Staine, K., Gochfeld, M., 1993. Fishing in contaminated waters: knowledge and risk perception of hazards by fishermen in New York City. *J. Toxicol. Environ. Health* 39, 95–105.
- Burger, J., Pflugh, K.K., Lurig, L., von Hagen, L., von Hagen, S.A., 1999a. Fishing in urban New Jersey: ethnicity affects information sources, perception, and compliance. *Risk Anal.* 19, 217–229.
- Burger, J., Stephens, W., Boring, C.S., Kuklinski, M., Gibbons, J.W., Gochfeld, M., 1999b. Ethnicity and risk: fishing and consumption in people fishing along the Savannah River. *Risk Anal.* 19, 427–438.
- Burger, J., Gaines, K.F., Gochfeld, M., 2001. Ethnic differences in risk from mercury among Savannah River fishermen. *Risk Anal.* 21, 533–544.
- Canadian Food Inspection Agency, 2002. Food safety facts on mercury and fish consumption. Available at: <http://inspection.gc.ca/english/corppaffr/foodfacts/mercurye.shtml>; accessed 17 August 2003.
- Carrington, C.D., Bolger, P.M., 2002. An exposure assessment for methylmercury from seafood for consumers in the United States. *Risk Anal.* 22, 689–699.
- Carrington, C.D., Cramer, G.M., Bolger, P.M., 1997. A risk assessment for methylmercury in tuna. *Water Air Soil Pollut.* 97, 273–283.
- Collings, S.E., Johnson, M.S., Leah, R.T., 1996. Metal contamination of angler-caught fish from the Mersey Estuary. *Marine Environ. Res.* 41, 281–297.
- Consumer Reports, 2001. America's fish: fair or foul? Available at: <http://www.consumerreports.org/special/consumerInteret/Reports/0102fis0.html>.
- Davidson, P.W., Myers, G.J., Cox, C., Axtell, C., Shamlaye, C., Sloane-Reeves, J., Cernichiari, E., Needham, L., Choi, A., Wang, Y., Berlin, M., Clarkson, T.W., 1998. Effects of prenatal and postnatal methylmercury exposure from fish consumption on neurodevelopment: outcomes at 66 months of age in the seychelles child development study. *JAMA* 280, 701–707.
- Denton, G.R.W., Burdon-Jones, C., 1996. Trace metals in fish from the great barrier reef. *Mar. Pollut. Bull.* 17, 201–209.
- Dey, P.M., Gochfeld, M., Reuhl, K.R., 1999. Developmental methylmercury administration alters cerebellar PSA-NCAM expression and Golgi sialyltransferase activity. *Brain Res.* 845, 139–151.
- DiFranco, J., Mower, B., 1994. Fish tissue contamination in the State of Maine—regional environmental monitoring and assessment program (REMAP). *Lake Reservoir Manage.* 9, 68–69.
- Ebert, E.S., 1996. Fish consumption and human health: developing partnerships between risk assessors and resource managers. *Am. Fisheries Soc. Symp.* 16, 261–270.
- Egeland, G.M., Middaugh, J.P., 1997. Balancing fish consumption benefits with mercury exposure. *Science* 278, 1904–1905.
- Environmental Protection Agency [EPA], 1998. Assessing human health risks from chemically contaminated fish and shellfish: a guidance manual. EPA-503/8-89-002, Appendix F. US Environmental Protection Agency, Cincinnati, Ohio.
- Food and Drug Administration [FDA], 1977. Food and Drug Administration Rules and Regulations, Part 161. Fish and Shellfish. Subpart B—Requirements for specific standardized fish and shellfish. *Federal Register* 47(50) 11,464–11,471.
- Food and Drug Administration [FDA], 1987. Compliance Policy Guide. Food and Drug Administration, Washington, D.C.
- Food and Drug Administration [FDA], 2001a. FDA consumer advisory. Available at: <http://www.cfsan.fda.gov/~dms/hgpdfloc.html>; Accessed 25 November 2003.
- Food and Drug Administration [FDA], 2001b. Mercury levels in seafood species. Available at: <http://www.cfsan.fda.gov/~frf/sea-mehg.html>; accessed 25 November 2003.
- Gerstenberger, S.L., Pratt-Shelley, J., Beattie, M.S., Dellinger, J.A., 1993. Mercury concentrations of walleye (*Stizostedion vitreum vitreum*) in 34 Wisconsin lakes. *Bull. Environ. Contam. Toxicol.* 50, 612–617.

- Gorman, J., 2003. Does mercury matter? experts debate the big fish question. *New York Times* July 29, 2003, p. F5.
- Graff, R.D., Philbert, M.A., Lowndes, H.E., Reuhl, K.R., 1993. The effect of glutathione depletion on methyl mercury-induced microtubule disassembly in cultured embryonal carcinoma cells. *Toxicol. Appl. Pharmacol.* 120, 20–28.
- Grandjean, P., Weihe, P., White, R.F., Debes, F., 1998. Cognitive performance of children prenatally exposed to "safe" levels of methylmercury. *Environ. Res.* 77, 165–172.
- Guallar, E., Sanz-Gallardo, M.I., van't Veer, P., Bođe, P., Aro, A., Gomez-Aracena, J., Kark, J.D., Riemersma, R.A., Martin-Moreno, J.M., Kok, F.J., 2002. Heavy metals and myocardial infarction study group. Mercury, fish oils, and the risk of myocardial infarction. *N. Engl. J. Med.* 347, 1747–1754.
- Hall, E.T., 1974. Mercury in commercial canned sea food. *J. Assoc. Off. Anal. Chem.* 57, 1068–1073.
- Hightower, J., Moore, D., 2002. Mercury levels in high-end consumers of fish. *Environ. Health Perspect.* 111, 604–608.
- Horn, E., 1992. Toxics in seafood. *Tidal Exch.* 3, 6–7.
- Hunter, D.J., Kazda, I., Chockalingam, A., Fodor, J.G., 1988. Fish consumption and cardiovascular mortality in Canada: an inter-regional comparison. *Am. J. Prevent. Med.* 4, 5–11.
- Hylander, L.D., Silva, E.C., Oliveira, L.J., Silva, S.A., Kuntze, E.K., Silva, D.X., 1994. Mercury levels in alto pantanal: a screening study. *Ambio* 33, 478–484.
- Institute of Medicine [IOM], 1991. *Seafood Safety*. National Academy Press, Washington, DC.
- International Foundation for Conservation of Natural Resources [IFCNR], 2003. *King of All Tunas & His Regal Court*. Available at: <http://fisheries.ifcnr.com/article.cfm?NewsID=231>. Accessed 25 November, 2003.
- Jacobson, J.L., Jacobson, S.W., 1996. Intellectual impairment in children exposed to polychlorinated biphenyls in utero. *New Engl. J. Med.* 335, 783–789.
- Jacobson, J.L., Humphrey, H.E.B., Jacobson, S.W., Schwartz, S.L., Mullin, M.D., Welch, R., 1989. Determinants of polychlorinated biphenyls (PCBs), polybrominated biphenyls (PBBs), and dichlorodiphenyl trichloroethane (DDT) levels in the sera of young children. *Am. J. Publ. Health* 79, 1401–1404.
- Jacobson, J.L., Jacobson, S.W., Humphrey, J.B., 1990. Effects of in utero exposure to polychlorinated biphenyls and related contaminants on cognitive functioning in young children. *J. Pediatr.* 116, 38–45.
- Kamps, L.R., Carr, R., Miller, H., 1972. Total mercury–monomethylmercury content of several species of fish. *Bull. Environ. Contam. Toxicol.* 8, 273–279.
- Kimbrough, R.D., 1991. Consumption of fish: benefits and perceived risks. *J. Toxicol. Environ. Health* 33, 81–91.
- Lacerda, L.D., Bidone, E.D., Giumaraes, A.F., Pfeiffer, W.C., 1994. Mercury concentrations in fish from the Itacaiunas–Parauapebas River system. Carajas region. Amazon Ann. Acad. Bras. Sci. 66, 373–379.
- Lange, T.R., Royals, H.E., Connor, L.L., 1994. Mercury accumulation in largemouth bass (*Micropterus salmoides*) in a Florida lake. *Archiv. Environ. Contam. Toxicol.* 27, 466–471.
- Mahaffey, K.R., 1999. Methylmercury: a new look at the risk. *Public Health Rep.* 114, 397–413.
- May, H., Burger, J., 1996. Fishing in a polluted estuary: fishing behavior, fish consumption, and potential risk. *Risk Anal.* 16, 459–471.
- Minnesota Department of Health [MDH], 1997. *An expectant mother's guide to eating Minnesota fish*. MDH, St. Paul, Minnesota.
- Myers, G.J., Davidson, P.W., Cox, C., Shamlaye, C.F., Palumbo, D., Cernichiar, E., Sloane-Reeves, J., Wilding, G.E., Kost, J., Huang, L.S., Clarkson, T.W., 2003. Prenatal methylmercury exposure from ocean fish consumption in the seychelles child development study. *Lancet* 361, 1686–1692.
- Nakagawa, R., Yumita, Y., Hiromoto, M., 1997. Total mercury intake from fish and shellfish by Japanese people. *Chemosphere* 35, 2909–2913.
- National Marine Fisheries Service [NMFS], 1989. *Fisheries of the United States 1988*. National Marine Fisheries Service, Current Fishery Statistics No. 8800.
- National Oceanic Atmospheric Administration [NOAA], 2002. Press release: Shrimp overtakes canned tuna as top US seafood. Available at: <http://www.publicaffairs.noaa.gov/releases2002/aug02/noaa02113.html>; accessed 17 August 2003.
- National Research Council, 2000. *Toxicological Effects of Methylmercury*. National Academy Press, Washington, DC.
- New Jersey Mercury Task Force, 2002. *Final Report of the New Jersey Mercury Task Force*. New Jersey Department of Environmental Protection, Trenton.
- Olsen, S.F., Secher, N.J., 2002. Low consumption of seafood in early pregnancy as a risk factor for preterm delivery: prospective cohort study. *Brit. Med. J.* 324, 447–454.
- Park, J.G., Curtis, L.R., 1997. Mercury distribution in sediments and bioaccumulation by fish in two oregon reservoirs: point-source and nonpoint-source impacted systems. *Arch. Environ. Contam. Toxicol.* 33, 423–429.
- Parvaneh, V., 1979. An investigation on the mercury contamination of Persian Guif fish. *Bull. Environ. Contam. Toxicol.* 23, 357–359.
- Plessi, M., Bertelli, D., Monzani, A., 2001. Mercury and selenium content in selected seafood. *J. Food Composition Anal.* 14, 461–467.
- Raines, B., 2003. *Your Deadly Diet*. Health (June 2003), 120–128.
- Reinert, R.E., Knuth, B.A., Kamrin, M.A., Stober, Q.J., 1991. Risk assessment, risk management, and fish consumption advisories in the United States. *Fisheries* 16, 5–12.
- Safina, C., 1998. *Song for a Blue Ocean*. Henry Holt, New York.
- Schantz, S.L., 1996. Developmental neurotoxicity of PCBs in humans: what do we know and where do we go from here? *Neurotoxicol. Teratol.* 18, 217–227.
- Schober, S.E., Sinks, T.H., Jones, R.L., Bolger, M.P., McDowell, M., Osterloh, J., Garrett, E.S., Canady, R.A., Dillon, C.F., Sun, Y., Joseph, C.B., Mahaffey, K.R., 2003. Bloodmercury levels in US children and women of childbearing age, 1999–2000. *JAMA* 289, 1667–1674.
- Sparks, P., Shepherd, R., 1994. Public perceptions of the potential hazards associated with food production: an empirical study. *Risk Anal.* 14, 799–808.
- Statistical Analysis System [SAS], 1995. *SAS Users' Guide*. Statistical Institute, Inc. Cary, NC.
- Stern, A.H., 1993. Re-evaluation of the reference dose for methylmercury and assessment of current exposure levels. *Risk Anal.* 13, 355–364.
- Stern, A.H., Korn, L.R., Ruppel, B.E., 1996. Estimation of fish consumption and methylmercury intake in the New Jersey population. *J. Expos. Anal. Environ. Epidemiol.* 6, 503–525.
- Stern, A.H., Gochfeld, M., Weisel, C., Burger, J., 2001. Mercury and methylmercury exposure in the New Jersey pregnant population. *Arch. Environ. Health.* 56, 4–10.
- Storelli, M.M., Stuffer, R.G., Marcotrigiano, G.O., 2002. Total and methylmercury residues in tuna-fish from the Mediterranean Sea. *Food Additiv. Contam.* 19, 715–720.
- Toth Jr., J.F., Brown, R.B., 1997. Racial and gender meanings of why people participate in recreational fishing. *Leisure Sci.* 19, 129–146.
- Veficer, C.M., Knuth, B.A., 1994. Communicating contaminant risks from sport-caught fish: the importance of target audience assessment. *Risk Anal.* 14, 833–841.

- Voegborlo, R.B., El-Methnani, A.M., Abedin, M.Z., 1999. Mercury, cadmium and lead content of canned tuna fish. *Food Chem.* 67, 341–345.
- Weihe, P., Grandjean, P., Debes, F., White, R., 1996. Health implications for Faroe islanders of heavy metals and PCBs from pilot whales. *Sci. Total Environ.* 186, 141–148.
- Yallouz, A., Campos, R.C., Louzada, A., 2001. Níveis de mercúrio em atum sólido enlatado comercializado na cidade do Rio de Janeiro. *Ciencias Tecnol. Aliment (Campinas)* 21, 1–4.
- Yess, N.J., 1993. US Food and Drug Administration survey of methylmercury in canned tuna. *J. AOAC Int.* 76, 36–38.



- Home
- News
- Main Categories
- Top News
- Nation
- States
- Washington/Politics
- World
- Election 2002
- Editorial/Opinion
- Health & Science
- Census
- Offbeat
- More News
- Columnists
- Lotteries
- City Guides
- Government Guide
- Talk Today
- Money
- Sports
- Life
- Tech
- Weather

Search

Site Web
By

SEARCH FOR NEWSPAPER ARTICLES
CLICK HERE

GET NEWS IN YOUR INBOX
Click here to get the Daily Briefing in your inbox

News



• [E-MAIL THIS](#) • [PRINT THIS](#) • [SAVE THIS](#) • [MOST POPULAR](#) • [SUBSCRIBE](#)

11/04/2002 - Updated 11:52 PM ET

If you eat a lot of fish, you may run health risk

By Anita Manning, USA TODAY

Health-conscious Americans have been told repeatedly that fish is good for the heart and the waistline. But there is growing concern that some seafood lovers are consuming high doses of mercury along with their fish dishes and could be suffering health problems as a result.

Women of childbearing age have long been warned to limit their fish intake to reduce the risk of exposing an unborn baby to mercury. But in a new study, a San Francisco physician says she discovered high levels of toxic mercury, called methylmercury, in blood and hair samples taken from dozens of her patients — men, women and children.

How contamination spreads

- The most common sources of mercury in air are coal-burning power plants, municipal waste combustors, medical waste incinerators and hazardous waste combustors.
- Mercury can contaminate water or land through the discharge of industrial wastewaters.
- Tiny particles of mercury travel through smokestacks into the air. They then fall onto soil or water.
- Mercury can accumulate in fish and wildlife. Small fish are eaten by big fish and fish-eating birds generally have higher levels of contamination.

Source: EPA.

Hightower, whose yearlong study of patients in her Bay

Many were suffering symptoms associated with low-level mercury poisoning, including hair loss, fatigue, depression, difficulty concentrating and headaches. The implication, she says, is that anyone who consumes a lot of fish, especially large steak fish such as swordfish and shark, could be at risk.

"You have people who have been told to eat fish because it's healthful, but they have not been told it contains contaminants," says physician Jane

RELATED STORIES

Latest news



- [Sharon, Netan U.S. peace pl](#)
- [Suicide bomb mall](#)
- [Israel reporter war efforts](#)
- [Netanyahu m: Cabinet](#)

Background



- [Developments](#)

BEYOND WORDS

Audio



- [USA TODAY: Israel's secret](#)

Graphics and galleries



- [Map: The Mid](#)

Area practice was published Friday in *Environmental Health Perspectives*, an online journal of the National Institute of Environmental Health Services, part of the National Institutes of Health.

Concern is widespread:

A Food and Drug Administration advisory committee recommended in July that the agency do research to assess the risks to women and young children who eat canned tuna. The amount of methylmercury per can is generally low, about 0.17 parts per million, but it can vary widely, says Michael Bender of the Mercury Policy Project, an advocacy group.

"Tuna is the most consumed fish in the country," Bender says. "If you're a pregnant woman and you eat over two cans of tuna per week, you can go over" safe levels of mercury. The FDA currently recommends that women who are or could become pregnant limit all fish to 12 ounces a week.

A survey of Hong Kong high school students found that as many as 10% eat enough fish to exceed safety limits for mercury exposure. The report, which prompted a Chinese government warning about consumption of shark and other large fish, found that the students' diets gave them a mercury exposure of 6.41 micrograms per kilogram (2.2 pounds) of body weight a week. The World Health Organization recommends a 5-microgram limit.

In September, the United Nations Environment Programme hosted a meeting in Geneva about ways to reduce mercury emissions around the world. A report from that meeting will be considered by environment ministers at a meeting of UNEP's governing council in February and could lead to consideration of an international treaty on mercury emissions.

Hightower's study and similar reports from other researchers who attended a recent meeting in Vermont, sponsored by the EPA, suggest that consumers who eat expensive fish are increasingly putting themselves at risk for mercury poisoning.

How much is safe?

The amount of canned tuna* that is safe to eat each week should be based on body weight:

- 25 lbs. 1 tablespoon.
- 50 lbs. 2 oz.
- 75 lbs. 3 oz.

"They are switching to fish to improve their health," says Caroline Smith DeWaal of the Center for Science in the Public Interest, but "they're being exposed to dangerously high levels of methylmercury." That's especially troubling if the consumers are women who plan to have children, says DeWaal, author

USA
TODAY
2001

All
the
News
to
your
inbox

Click
to sign up!

- * 100 lbs. 5 oz.
- * 125 lbs. 1 can or 6 oz.
- * 150 lbs. 8 oz.
- * 175 lbs. 9 oz.
- * 200 lbs. 10 oz.

* Choose albacore, chunk or chunk light varieties, which have less mercury than solid white or chunk white types.

Source: *Fish Facts for Good Health*, publication of the Washington Department of Health.

of the recently published *Is Our Food Safe?* "It is critical that women of childbearing age stop eating this fish from six months to a year before becoming pregnant."

The Centers for Disease Control and Prevention estimates about 8% of women in this age group have enough mercury in their bodies to pose a risk of having babies with mild learning problems.

No set standards

Mercury released from power plants, municipal waste facilities and medical incinerators is the primary source of methylmercury in fish. Methylmercury is an organic form of mercury that is different from what is in mercury thermometers or what goes up smokestacks when coal is burned.

Mercury is converted to methylmercury by bacteria in water. So when people are talking about mercury in fish, they are really talking about the toxic methylmercury. What makes it dangerous to health is that it is hard for the body to eliminate, so it can build up and may affect the nervous system. Most human exposure to methylmercury is through fish consumption.

The FDA ceased its large methylmercury sampling program in 1998, and today federal agencies conduct only limited testing of fish for methylmercury. The industry does, too, on a voluntary basis, says Rhona Applebaum, a scientist with the National Food Processors Association. "Whether it's mercury or any other defect, chemical or microbial, the industry does regular testing" to assure that the product meets FDA standards.

"We do know tuna contains methylmercury," she says, but mercury is "naturally occurring, so on a daily basis people are exposed. It's not at levels that will result in acute toxicity unless people are not practicing basic tenets of nutrition: balance, variety and moderation."

Studies show women ages 15-44 eat canned tuna 1.5 times a month, well within the range of safety, but too much of anything can be harmful, she says. "If people are going to consume one type of food literally ad nauseam, there's going to be an impact."

The FDA and Environmental Protection Agency differ on what they consider acceptable levels and measure it differently. The FDA, which regulates commercially caught fish, sets an "action level" of 1 part per million. If higher levels are reported, the FDA can remove the fish from the market, though critics say that rarely occurs. The EPA has a "reference dose" that says people can be exposed to .1 microgram per kilogram of

body weight per day, which is roughly 5 to 7 micrograms per day for someone who weighs 100 to 154 pounds, says Kate Mahaffey of the EPA. That's about a fifth of the amount the FDA considers safe.

The FDA's standard permits about 480 micrograms of methylmercury in one pound of fish, she says. "If fish is that contaminated, and you're trying to keep in the 5 to 7 micrograms per day range, you can't eat much of that fish."

But the EPA does not regulate commercial fish. It works with state environmental and health departments to test local rivers and other bodies of water where recreational fishing is done and where mercury levels may be high because of local pollution. When high levels of mercury are detected in the water, the states post fish advisories to warn consumers.

"There are fish advisories in most states for mercury," says Michael Gochfeld, professor of environmental medicine at the Robert Wood Johnson Medical School in Piscataway, N.J.

Even in areas where industrial pollution has been reduced, the problem persists because of atmospheric pollution drifting from other areas, he says. "Virtually all our mercury-polluting industry in New Jersey is closed," he says, but state health officials regularly warn residents not to eat fish from specific lakes and rivers where mercury levels are high. Ten states also warn pregnant women to limit consumption of canned tuna and other commercial seafood.

Yet health experts point out that fish is an important part of a balanced diet. It's full of vitamins and other nutrients, including omega-3 fatty acids, which help lower cholesterol and blood pressure and reduce the risk of cardiac arrest. And it's low in calories.

What fish should you eat?

Fish that have high levels of mercury:

- Swordfish
- Shark
- Tilefish
- King mackerel
- Tuna (steak)

Fish that have generally low levels of mercury:

- Salmon
- Flounder
- Cod
- Catfish
- Trout
- Pollock
- Clams
- Shrimp
- Scallops
- Lobster

Source: *Fish Facts for Good Health*, publication of the Washington Department of Health.

How that squares with mercury poisoning is "a very difficult message to convey," says epidemiologist Tom Sinks of the National Center for Environmental Health, part of the CDC. "Fish is a vehicle by which people are exposed to mercury. But at the same time, fish is a good source of protein and nutrients, an important part of the diet, and one we want people to eat in a healthy way."

He says the fish that are high in omega-3, such as salmon and sardines, are low on the mercury scale. "We want to encourage people not to avoid fish, but to advise them that some fish have higher levels of mercury, and if they're concerned, they should avoid those fish," he says.

The trouble, Hightower says, is that some people appear to be more sensitive to methylmercury than others. The EPA and the National Academy of Sciences recommend keeping mercury levels in blood at 5 micrograms per liter or less. In Hightower's study, patients' blood levels ranged from 2 to nearly 90 micrograms per liter. Symptoms varied widely and did not always correlate with the burden of methylmercury.

"There were some with elevated levels who had no symptoms. There are some with low levels with symptoms," she says. "It is unclear whether these patients are having symptoms due to direct effects of mercury or a reaction to it," she says. But, she adds, most people can withstand a bee sting, while others go into shock. "We recognize there are severe reactions to very minuscule quantities of certain agents."

Hightower says it's not known how many people might be affected by methylmercury, and she can't prove that the symptoms her patients suffered were caused by overconsumption of fish, but "the funny thing is, people got better when they stopped eating it."

Not eating fish helped

That's what happened to Wendy Moro, 40, a marketing consultant who lives with her husband and son in a suburb of San Francisco. Until April 2001, she says, she was the picture of health. A 110-pound bundle of energy, she ran several miles a day, danced ballet, lifted weights. She also ate fish two to five times a week, at home and at the Bay Area's better restaurants.

"On the West Coast, we eat a lot of fish," she says. "It's an affluent community, and fish is accessible and popular. You go out for dinner. People don't go out for T-bone steaks anymore. It's all fish."

She ate tuna for lunch a couple of times a week, and the family would have seafood for dinner regularly, often choosing steak fish such as ahi tuna or halibut. "We just looked for what was fresh," she says. "I thought I was being really healthy, not eating meat, eating lots of fish."

The first sign of trouble was severe fatigue — "the kind where it is impossible to stay awake for more than a few hours at a time," she says. Then pain and weakness in her limbs worsened to the point where she could barely stand. A series of doctors diagnosed or tested her for multiple sclerosis, muscular dystrophy, chronic fatigue syndrome, mononucleosis, diabetes insipidus. One suggested she be evaluated for mental illness.

Finally, she was referred to Hightower, who tested her for mercury poisoning. Moro's blood level was 17, more than three times the recommended level, though still below what some doctors think is enough to cause such severe symptoms.

When Moro stopped eating fish, her symptoms began to disappear. Now, she says, she's "about 85%" back to normal. She keeps a file on mercury that she gives to friends who are thinking about having a baby.

If it could happen to her, it could happen to anyone, she says. "I'm such an average Jane. I live in a suburb; I have 1.5 kids, if you count my dog. I'm not a super-fanatic, not a triathlete. I'm not super-rich or poor. I'm just an average Joe-USA TODAY. That's what's scary."

Alan Stern, chief of the New Jersey Department of Environmental Protection who served on a National Academy of Sciences committee on methylmercury two years ago, says it's too soon to draw firm conclusions from Hightower's study. "I would consider it to be the very early stages of a clinical case description, and it's not at a point yet where it can be translated into a public health message," he says.

Such reports, he says, "call our attention to the potential of health effects at low levels of exposure (to methylmercury), but they don't make an open-and-shut case."

Even the relief of symptoms reported by people who stop eating fish is inconclusive, he says, because it is "hard to distinguish that from a placebo effect. From an objective standpoint, one cannot say this association goes to the next step of cause and effect."

But if nothing else, Stern says, consumers and doctors should be alert to the possibility that small exposures to mercury in fish might cause symptoms. His cautionary conclusion: "Individuals should choose their diets wisely."

[Front Page](#) [News](#) [Money](#) [Sports](#) [Life](#) [Tech](#) [Weather](#) [Marketplace](#)
[USA TODAY Travel](#) [USA TODAY Careers Network](#)
[Terms of service](#) [Privacy Policy](#) [How to advertise](#) [About us](#)
 © Copyright 2002 USA TODAY, a division of [Gannett Co. Inc.](#)